

Evolution of introduced Chinook salmon (*Oncorhynchus tshawytscha*) in Lake Huron: emergence of population genetic structure in less than 10 generations

Ho Y. Suk*, Bryan D. Neff, Kevin Quach, Yolanda E. Morbey

Department of Biology, University of Western Ontario, London, ON, Canada

Accepted for publication October 21, 2011

Abstract – Population genetic structure was detected in Chinook salmon *Oncorhynchus tshawytscha* in their non-native range of Lake Huron using microsatellite DNA. All Chinook salmon in this system descend from Green River, Washington cohorts, originally transplanted to Michigan hatcheries in the late 1960s. We tested for population genetic differentiation of age 0 fish collected from 13 rivers and two hatcheries in 2007. The amount of genetic differentiation among collection sites was low but statistically significant, with F_{ST} values ranging from 0.036 to 0.133 and R_{ST} values ranging from 0.008 to 0.157 for specific loci. Based on pairwise F_{ST} and R_{ST} values and Bayesian cluster analysis, the Maitland River population in the Main Basin of Lake Huron was genetically distinct from the remaining collection sites. Based on analysis of bycatch data from commercial gill net fisheries, Chinook salmon likely colonised the Main Basin by 1975 (10 generations ago) and the North Channel and southern Georgian Bay regions by 1980 (eight generations ago). Thus, population genetic structure has emerged in Lake Huron Chinook salmon in <10 generations.

Key words: introduced species; Salmonidae; population genetics; evolutionary divergence; rapid evolution

Introduction

Chinook salmon (*Oncorhynchus tshawytscha* Walbaum) in Lake Huron provide an exceptional opportunity to explore the dynamics of contemporary evolution at an early postcolonisation phase, in a large number of populations, and at a large spatial scale. In the 1960s, the lower four Laurentian Great Lakes were suffering the consequences of the extirpation of almost all populations of lake trout (*Salvelinus namaycush* Walbaum) from overharvesting and predation by sea lamprey (*Petromyzon marinus* Linnaeus) (Coble et al. 1990; Eshenroder et al. 1992). In response, fisheries managers initiated a Pacific salmonid stocking programme to control burgeoning alewife

(*Alosa pseudoharengus* Wilson) populations and provide recreational fishing opportunities to coastal communities (Hansen 1999; Crawford 2001). This programme has involved artificial breeding and the release of hatchery-reared juveniles (age 0) into tributaries and along lakeshores. Since the initiation of the stocking programme, Chinook salmon have established feral populations¹ in Lake Michigan, Lake Superior, Lake Huron, and, to a lesser extent, Lake Ontario (e.g., Peck et al. 1999; Weeder et al. 2005; Smith et al. 2006; Johnson et al. 2010). Indeed, large-scale mark-recapture studies show that wild Chinook salmon can comprise the majority of the recreational catch in some regions (Peck et al. 1999; Johnson et al. 2010). In Lake Huron, Chinook salmon are known to

Correspondence: Y. E. Morbey, Department of Biology, University of Western Ontario, 1151 Richmond Street N, London, ON N6A 5B7, Canada. E-mail: ymorbey@uwo.ca

*Present address: Department of Biology, Yeungnam University, 214-1 Daedong, Gyeongsan City, Gyeongsangbuk-do 712-749, Korea.

¹Defined as a group of individuals using a particular river for breeding and, due to natal homing, unlikely to interbreed with other such groups.

reproduce in at least 17 tributaries, with most of these associated with the Niagara Escarpment regions of Ontario and the upper peninsula of Michigan (Marklevitz et al. 2011). Little is yet known about the degree of genetic differentiation among these feral populations of Chinook salmon and whether they are sufficiently isolated to be undergoing local adaptation.

The stocking programme for Chinook salmon in Lakes Michigan and Huron began in 1966–1968 with the transfer of embryos of Green River, Washington, origin to the state of Michigan (Weeder et al. 2005). Age 0 fish were released into tributaries of Lake Michigan starting in 1967, and to a lesser extent, Lake Huron starting in 1968 (Great Lakes Fishery Commission [GLFC] Fish Stocking database; Crawford 2001). A smaller transfer of Green River origin embryos from the University of Washington hatchery occurred in 1967–1969 to Alpena Community College with subsequent release of age 0 fish into the Thunder Bay River (J. Johnson, Michigan Department of Natural Resources, personal communication). Since then, no additional egg transfers from the west coast have occurred. Currently, seven hatcheries raise age 0

fish for release into Lake Huron: three state-operated facilities in Michigan obtain gametes from adults returning to the Swan River, and four community-run facilities in Ontario obtain gametes from adults returning to local tributaries of Lake Huron (primarily the Sydenham River, Saugeen River, Beaver River, and sites on Manitoulin Island; Fig. 1).

The first evidence of successful natural reproduction of Chinook salmon in Lake Huron was in 1983, when wild age 0 fish were captured in the Sydenham and nearby Pottawatomi Rivers (Kerr & Perron 1986). This record was followed in 1985–1987 by the discovery of age 0 out-migrants during weir surveys in these and four additional tributaries in the southern Georgian Bay region (Beaver, Bighead, Sauble, and Saugeen Rivers; Fig. 1) (Kerr 1987). In addition to the evidence of spawning in rivers, lake spawning was observed in the North Channel of Lake Huron in 1987 (Powell & Miller 1990). Given that hatcheries were not operating in Ontario until 1984, the early colonisers of Ontario tributaries (i.e., ‘strays’) were most certainly hatchery-reared fish from Lake Michigan or Michigan waters of Lake Huron (Kerr 2006).




Fig. 1. Map of Lake Huron showing the Ontario Ministry of Natural Resources fisheries assessment areas (numbers 01–25), collection sites for Chinook salmon (filled circles) and other landmarks (filled triangles). Publicly available layers were used to make the maps in ArcMap 9.2 (ESRI, Redlands, CA, USA). The maps are projected using the North American Albers Equal-Area Conic Projection. Site codes are defined in Table 2.

Typically, adults from wild populations show philopatry to natal spawning areas because of imprinting before and during the parr-smolt transformation (Dittman & Quinn 1996; Hendry et al. 2004). The release of hatchery-reared fish at non-natal sites disrupts imprinting and subsequent homing ability (e.g., Quinn 1993; Candy & Beacham 2000) and may have promoted colonisation of Ontario tributaries of Lake Huron.

Our objective was to test for population genetic structure of Chinook salmon in Lake Huron using microsatellite loci. Previously, Weeder et al. (2005) observed a lack of genetic differentiation among feral populations of Chinook salmon in Lake Michigan based on data from allozyme loci. This was attributed to the combined effects of recent colonisation and straying by hatchery-reared fish. However, microsatellites may be better loci for discriminating population structure because of their relatively high mutation rate. To complement the genetic results, we used commercial fisheries monitoring data to establish when Chinook salmon colonised different regions of Lake Huron and how many generations had passed since colonisation. We hypothesised that some feral populations of Chinook salmon would be genetically distinct because of founder effects, postcolonisation genetic drift and/or natural selection.

Methods

Collection of age 0 Chinook salmon

Age 0 Chinook salmon of wild and hatchery origin were collected from 15 collection sites within the Lake Huron watershed during April–June 2007 (Fig. 1). Wild fish, identified by their small size and lack of any fin clips, were collected from 13 feral populations, whereas hatchery-origin fish were collected from two hatcheries. Our fish samples comprised a subset of those used in a study of otolith microchemistry from 17 known feral populations and seven hatcheries in Lake Huron (Marklevitz et al. 2011). Because our objective was to assess genetic differentiation among extant populations in Lake Huron, not divergence from the source population, we did not collect archived samples from the Green River. Previous research already indicates a significant founder effect because of the transfer of gametes from the Green River to Michigan (Weeder et al. 2005). Wild fish were sampled in rivers by electrofishing suitable juvenile habitats (e.g., side channels of main rivers) near areas where spawning had been observed by local biologists. In these systems, finding juveniles was challenging and took a lot of effort. Hatchery fish were sampled from raceways using dip nets. Fish were euthanised with clove oil or MS222. From each

individual fish ($n = 20\text{--}24$ per collection site), a tail fin clip was preserved in 95% ethanol.

The feral populations in our study have variable stocking histories, with no recorded hatchery releases in the Root River, Spanish River, Manitou River, Beaver River and Bighead River, <200,000 hatchery releases in the Nottawasaga River ($n = 60,000$ in 1988), Maitland River ($n = 100,000$ in 1985 and 1987) and Carp River ($n = 125,114$ from 1980 to 1985), 0.5–1 million hatchery releases in the Sauble River ($n = 712,776$ from 1986 to 2002) and Saugeen River ($n = 860,798$ from 1985 to 1998), and over 1 million hatchery releases in the St Marys River ($n = 1,730,866$ from 1977 to 2003), Sydenham River ($n = 3,612,675$ from 1985 to 2002) and Nunn's Creek ($n = 5,984,490$ from 1988 to 2010) (GLFC Fish Stocking Database). The two hatcheries in our study collect gametes from adults returning to local tributaries, with the Lake Huron Fishing Club (LHFC) usually collecting from the Saugeen River and Sydenham River and the Sydenham Sportsmen's Association (SSA) usually collecting from the Sydenham River. The Beaver River also is a gamete collection site in some years. The LHFC releases fish at a variety of sites in local tributaries and along lakeshores, whereas the SSA releases the majority of its fish in the Sydenham River (GLFC Fish Stocking Database).

Microsatellite DNA genotyping

We used nine previously published microsatellite loci for genotyping: *Oneμ3*, *Oneμ8*, *Oneμ14* (Scribner et al. 1996), *Ssa85* (O'Reilly et al. 1998), *Omy207*, *Omy325* (O'Connell et al. 1997), *OtsG311*, *OtsG432* and *OtsG474* (Williamson et al. 2002). All forward primers were fluorescently labelled (Sigma-Genosys, The Woodlands, TX, USA). Genomic DNA was extracted from the tail fin of each fish by the Wizard Genomic DNA purification kit (Promega, Madison, WI, USA), following the manufacturer's protocol. We used a T1 Thermocycler (Biometra, Göttingen, Germany) to amplify the microsatellites with the following programme: 94 °C for 10 min, 35 cycles of 30 s denaturing at 94 °C, 30 s at 58 °C (*Oneμ3*, *Oneμ8*, *Oneμ14*, *Omy207*, *Omy325* and *OtsG474*) or 61 °C (*Ssa85*, *OtsG311* and *OtsG432*) for annealing the primers, and 30 s at 72 °C for nucleotide extension and final elongation at 72 °C for 10 min. Each 10-ml PCR contained 75 ng of total DNA, 3 mM MgCl₂, 10× PCR buffer (Invitrogen Life Technologies, Carlsbad, CA, USA), 0.25 mM of each deoxynucleotide (Sigma-Aldrich, St Louis, MO, USA), 0.25 units Taq DNA polymerase (Invitrogen Life Technologies) and 0.25 mM of each forward and reverse primer (Invitrogen Life Technologies). PCR products were

size-fragmented following the standard protocol for the CEQ 8000 Genetic Analysis System (Beckman Coulter, Mississauga, ON, Canada).

Genetic data analysis

Genetic diversity was quantified by the number of alleles per locus, observed heterozygosity H_O , and expected heterozygosity H_E (Nei 1987) using ARLEQUIN 3.1 (Schneider et al. 1997). Deviations from Hardy–Weinberg equilibrium (HWE) were tested for each locus–population combination using the GENEPOL 3.1d package (Raymond & Rousset 1995) with the exact test based on a Markov chain method (Guo & Thompson 1992). For tests over all loci, the significance of departures from expected HWE genotype frequencies was assessed (see F_{IS} in Table 1). MICRO-CHECKER (van Oosterhout et al. 2006) was used to detect possible technical artefacts, such as null alleles, large allele dropout and accidental scoring of stutter bands. ML-RELATE (Kalinowski et al. 2006) was applied to test whether there was a family effect in our sampling and analyses.

We used multiple approaches to examine the extent of genetic differentiation and structuring among collection sites. The statistics make different assumptions about the mode of mutation at microsatellites and are derived from either the infinite allele model (IAM) or the stepwise mutation model (SMM). Although statistics based on the SMM typically suffer high sampling variances, those based on the IAM can underestimate differentiation among highly structured populations in which mutation rates are greater than migration rates (Balloux & Lugon-Moulin 2002). As a conservative approach, both families of statistics were used in the present study. We estimated pairwise F_{ST} (Weir & Cockerham 1984) and pairwise R_{ST} (Michalakis & Excoffier 1996) values using GENEPOL. Multidimensional scaling analysis (MSA), based on pairwise F_{ST} values, was used to visualise the genetic relationships among collection sites using VISTA 5.6.3 (Young 1996). Differentiation by distance was anal-

ysed by calculating the correlation coefficient of $F_{ST}(1 - F_{ST})^{-1}$ and geographically shortest distance along the water between collection sites. The significance of the relationship was evaluated with 100,000 permutations in a Mantel test implemented in TFPGA v1.3 (Miller 1997).

Population genetic structure was first analysed in a Bayesian framework based on an individual-level clustering method. This approach can identify hidden population structure by clustering individuals into genetically distinguishable groups on the basis of allele frequencies and linkage disequilibrium without a priori assumptions of population structure (Corander et al. 2004). This analysis was performed in the BAPS 4.1 program (Corander et al. 2004). Three iterations were run for each value of K (the putative number of clusters) using the admixture model with Markov chain Monte Carlo (MCMC) values of 100,000 each, and values of $K = 1–30$. The number of significant clusters was considered to be the value of K corresponding to the plateau of log-likelihood ratios. Next, hierarchical genetic structuring was analysed by assessing the relative contributions of among-group (based on the outcome of the Bayesian cluster analysis), among-population and within-population components using an analysis of molecular variance (AMOVA; Excoffier et al. 1992) using the program ARLEQUIN 2.0 (Schneider et al. 1997). The AMOVA was carried out with both IAM- and SMM-based distance measures, and the significance of each hierarchical level was tested with 1000 permutations.

Sampling of commercial fisheries

Commercial fisheries monitoring data (1975–2008) were used to estimate the generation time of Chinook salmon in Lake Huron and to determine when the three main regions (Main Basin, Georgian Bay and North Channel; Fig. 1) were colonised. In combination with the genetic data, these data provide information on the time course for genetic differentiation. In Lake Huron, on-vessel catch samplers collect

Table 1. Summary of the nine microsatellite loci used to study the genetic differentiation of Chinook salmon among collection sites in Lake Huron.

Loci	Size range (bp)	A	H_O	H_E	F_{IS}	F_{ST}	R_{ST}
<i>Omy325</i>	81–103	13	0.870	0.844	-0.095	0.066*	0.044*
<i>One3</i>	161–169	4	0.603	0.469	-0.343	0.044*	0.044*
<i>OtsG474</i>	150–206	13	0.773	0.726	-0.088	0.036*	0.013*
<i>Ssa85</i>	118–168	17	0.866	0.851	-0.052	0.043*	0.038*
<i>OtsG311</i>	230–362	32	0.928	0.951	-0.028	0.053*	0.035*
<i>Omy207</i>	77–81	4	0.092	0.081	-0.181	0.133*	0.157*
<i>OtsG432</i>	105–184	14	0.875	0.860	-0.067	0.051*	0.008*
<i>One8</i>	139–195	12	0.773	0.734	-0.093	0.055*	0.017*
<i>One14</i>	188–244	15	0.638	0.848	0.157*	0.102*	0.034*

Data include allele size range in base pairs (bp), total number of alleles (A), mean observed (H_O) and expected heterozygosity (H_E), F_{IS} , F_{ST} and R_{ST} . * $P < 0.05$.

standard data on fisheries catch and bycatch (e.g., Gile & Milne 2006). Fishing effort is targeted primarily towards lake whitefish (*Coregonus clupeaformis* Mitchell) using trap nets or gill nets of variable mesh size. Chinook salmon caught as bycatch are enumerated, measured and in some cases aged using scales. We estimated mean generation time based on data from fish assigned as mature (will spawn in the current year), excluding those ($n = 4$) with assigned ages of 0. Annual bycatch of Chinook salmon was estimated for the gill net fishery as catch rate (number captured·km gill net $^{-1}$ ·24 h $^{-1}$). In an effort to account for the potential effect of mesh size on catch rates, only sets using large mesh (>90 mm stretched mesh) were used in the determination of catch rates. Median annual catch rates were calculated for each 5-year period in each fisheries assessment area (Fig. 1).

Results

Observed heterozygosity (H_O) was high (average H_O 's > 0.6) for all loci except for *Omy207* with a $H_O = 0.09$ (Table 1). The number of alleles per locus ranged from 4 to 32. Based on its high F_{IS} value, only *Oneμ14* had genotype frequencies that departed significantly from HWE within populations. MICRO-CHECKER revealed that *Oneμ14* had a significant probability of null alleles in five populations (Table 2). There was no significant departure from HWE at the population level (Table 2). Mean F_{ST} values for specific loci ranged from 0.04 to 0.13, whereas mean R_{ST} values ranged from 0.01 to 0.16. Genetic differentiation among populations was evident based on significant values of F_{ST} and R_{ST} for all loci (Table 1). Mean F_{ST} across all loci was 0.065, indicating that $n = 20\text{--}44$ per population warranted further statistical

analyses (Kalinowski 2005). Exact tests for genetic linkage disequilibrium did not reveal any significant effects, suggesting the absence of physical linkage of the loci (data not shown). Maximum likelihood estimation of relatedness and relationship by ML-RELATE did not reveal any family effect in our sampling and genetic analysis; more than 90% of the possible individual pairs were unrelated in all collection sites (95% confidence set with 10,000 randomisations).

Based on pairwise F_{ST} values averaged across loci, differentiation was strongest in the Maitland River population ($F_{ST} = 0.09\text{--}0.21$, 14 significant F_{ST} values), followed by the Root River population ($F_{ST} = 0.03\text{--}0.14$, 12 significant F_{ST} values), Sauble River population ($F_{ST} = 0.01\text{--}0.11$, 8 significant F_{ST} values), Nunn's Creek population ($F_{ST} = 0.05\text{--}0.21$, 5 significant F_{ST} values) and the LHFC Hatchery ($F_{ST} = 0.02\text{--}0.16$, 5 significant F_{ST} values; Table 3). Based on pairwise R_{ST} values averaged across loci, differentiation was most apparent for the Maitland River population (14 significant R_{ST} values; Table 3). The distinct nature of the Maitland River population was also evident from the multidimensional scaling plot of pairwise F_{ST} values (Fig. 2). The Mantel test failed to find any correlation between genetic differentiation and geographic distance (pairwise F_{ST} values: $r = 0.164$, $P = 0.98$; pairwise R_{ST} values: $r = 0.021$, $P = 0.54$).

In the Bayesian analysis of population structure, log-likelihood ratios reached a plateau at $K = 6$, indicating that most of the genetic variation was captured in six genetic clusters (Table 4). In 12 collection sites, most individuals (43–100%) were assigned to cluster 3. In contrast, the majority of individuals from Nunn's Creek (75%), Maitland River (71%) and Saugeen River (50%) were assigned to

Table 2. Summary of genetic characteristics of Chinook salmon from 15 collection sites in Lake Huron.

Collection site	Code	n	AR	H_E	H_O	F_{IS}	Loci departing from HWE
Sydenham River	SY	44	5.610	0.658	0.675	-0.025	<i>Oneμ14</i>
Nunn's Creek	NN	20	4.392	0.569	0.667	-0.101	
Sauble River	SB	20	5.940	0.750	0.753	-0.004	<i>Oneμ14</i>
Saugeen River	SG	20	4.923	0.660	0.703	-0.064	
St Marys River	StM	20	5.741	0.707	0.722	-0.022	<i>Oneμ14</i>
Spanish River	SP	25	6.061	0.703	0.730	-0.040	
Nottawasaga River	NT	25	5.142	0.665	0.699	-0.052	
Manitou River	MN	22	5.850	0.705	0.756	-0.075	<i>Oneμ14</i>
Beaver River	BV	20	5.633	0.713	0.682	0.045	
SSA Hatchery	SSA	20	5.439	0.673	0.708	-0.053	
Root River	RT	20	5.244	0.710	0.792	-0.119	
LHFC Hatchery	LHFC	20	5.050	0.605	0.689	-0.144	
Carp River	CR	20	5.888	0.683	0.751	-0.104	
Maitland River	MT	21	3.997	0.562	0.644	-0.151	
Bighead River	BH	21	5.664	0.691	0.724	-0.049	<i>Oneμ14</i>

Shown are the collection site, collection site code, sample size (n), allelic richness (AR), mean expected (H_E) and observed heterozygosity (H_O), F_{IS} , and any loci departing from Hardy-Weinberg equilibrium.

SSA, Sydenham Sportsmen's Association; LHFC, Lake Huron Fishing Club.

Table 3. Genetic differentiation parameters (F_{ST} and R_{ST}) for pairwise comparisons of Chinook salmon from 15 collection sites in Lake Huron.

Site	SY	NN	SB	SG	StM	SP	NT	MN	BV	SSA	RT	LHFC	CR	MT	BH
SY	0.042	0.029	0.070	0.007	0.005	0.013	0.019	0.011	0.011	0.001	0.001	0.021	0.055*	-0.002	
NN	0.082	0.045	0.199	0.017	0.020	0.128	0.119	0.093*	0.028	0.044	0.017	0.108	0.022*	-0.003	
SB	0.047*	0.086	0.065	-0.002	0.036	0.052	0.023	-0.005	0.032	-0.010*	-0.019	0.009	0.059*	0.017	
SG	0.045	0.124	0.045*	0.076*	0.094	0.045	0.049	0.020	0.141	0.025*	0.070	0.025	0.180*	0.091	
StM	0.031	0.048	0.014	0.051	-0.003	0.062	0.054	0.022	0.017	-0.009	-0.012*	0.044	0.006*	-0.020*	
SP	0.017	0.033	0.028	0.048	0.003	0.042	0.041	0.022	-0.011	-0.007	-0.006	0.046	-0.004*	-0.013	
NT	0.026	0.100	0.037*	0.067	0.037	0.033	-0.022	0.000	0.043	0.013	0.024	-0.012	0.128*	0.053	
MN	0.033	0.074	0.029*	0.055	0.020	0.015	0.031	-0.014	0.030	0.004	0.008	-0.028	0.112*	0.046	
BV	0.030	0.101*	0.018	0.047	0.026	0.024	0.036	0.034*	0.029	-0.019	-0.019	-0.010	0.087*	0.033	
SSA	0.034	0.090	0.042	0.070	0.026	0.033	0.042	0.043	0.030	0.006	-0.010	0.040	0.018*	0.007	
RT	0.065*	0.076*	0.054*	0.074*	0.028*	0.029*	0.078*	0.050*	0.058	0.066*	-0.025	0.002	0.035*	0.004	
LHFC	0.052	0.067*	0.067*	0.102	0.048	0.023	0.083	0.054	0.057	0.067	0.082*	0.003	0.021*	-0.009	
CR	0.051	0.087	0.034	0.067	0.038	0.030	0.054	0.023	0.045	0.059	0.073*	0.058*	0.113*	-0.009	
MT	0.093*	0.205*	0.111*	0.116*	0.125*	0.117*	0.106*	0.127*	0.101*	0.118*	0.137*	0.162*	0.153*	0.049*	
BH	0.034	0.067	0.039*	0.054	0.023	0.013	0.053	0.039	0.032	0.044	0.035	0.054	0.053	0.106*	

Estimates of F_{ST} appear below the diagonal, and estimates of R_{ST} appear above the diagonal (site abbreviations defined in Table 2). * $P < 0.05$.




Fig. 2. Multidimensional scaling plot of pairwise F_{ST} at nine microsatellite loci, showing the genetic relationships among Chinook salmon collection sites. The circles encompass groupings with genetic similarity. Site codes are defined in Table 2.

different clusters (2, 4 and 6, respectively). Saugeen River also had a large fraction of individuals (40%) assigned to cluster 3. Based on the clustering pattern, we considered Nunn's Creek, Maitland River and the set of the remaining 13 collection sites as three separate groups in the AMOVA. Under the IAM, there was small but significant genetic variance among groups (1.9% of total variance, $P < 0.0001$) and among collection sites within groups (5.2%, $P < 0.0001$). Most of the genetic variance was found in the residual component (92.9%, $P = 0.065$). The same conclusions were obtained under the SMM. There was small but significant genetic variance among groups (1.3%, $P = 0.001$) and among collection sites with groups (2.7%, $P = 0.003$), and most of the genetic variance was found in the residual component (96.0%, $P = 0.2$).

Table 4. Summary of the Bayesian analysis of population structure (proportions in each cluster) of Chinook salmon in Lake Huron.

Collection site	Cluster					
	1	2	3	4	5	6
Sydenham River	0.05	0.68*	0.23	0.02	0.02	
Nunn's Creek	0.75*	0.25				
Sauble River	0.05	0.95*				
Saugeen River	0.10	0.40	0.50*			
St Marys River	0.25	0.70*			0.05	
Spanish River	0.28	0.64*	0.04	0.04		
Nottawasaga River	0.16	0.80*	0.04			
Manitou River	0.14	0.77*	0.05	0.05		
Beaver River	0.10	0.85*			0.05	
SSA Hatchery		1.00*				
Root River		0.95*			0.05	
LHFC Hatchery	0.05	0.89*			0.05	
Carp River	0.11		0.89*			
Maitland River	0.05	0.10	0.14			0.71*
Bighead River	0.05	0.43*	0.10	0.33	0.10	

*Dominant clusters for each population.

Maturity and age information was available for 716 of the 19,073 Chinook salmon caught as bycatch in Ontario waters of Lake Huron from 1975 to 2008. The mean age of the 216 mature individuals was 2.2 years for a mean generation time of 3.2 years. In 1975–1979, the majority of Chinook salmon in the large mesh gill net fishery was captured in the Main Basin (in particular, areas 5, 8 and 9; Fig. 3). By 1980–1984, Chinook salmon started to be caught in southern Georgian Bay (areas 19–20) and the North Channel (areas 21–24). Throughout the 1980s, catch rates were highest in areas 5 and 9 of the Main Basin. By 1990–1994, catch rates in area 20 of southern Georgian Bay peaked and exceeded those elsewhere. During 1995–2004, catch rates in the North Channel (in particular, areas 21–23) peaked and exceeded those elsewhere. By 1995, Chinook salmon were being caught in the majority of assessment areas in Lake Huron.




Fig. 3. Median catch rates ($\text{number captured} \cdot \text{km}^{-1} \cdot 24 \text{ h}^{-1}$) of Chinook salmon in large mesh, commercial gill net fisheries by assessment area within 5-year periods from 1975 to 2008. The xs indicate assessment areas without data, either because of no commercial fishing of large mesh gill nets or lack of catch samplers.

Discussion

Since their introduction in the 1960s, Chinook salmon have become widespread in Lake Huron at every life history stage (Johnson et al. 2010; Marklevitz et al. 2011) and have played a vital role in the lake's food web dynamics (Dobiesz et al. 2005). They also support a multi-million dollar recreational fishing industry in Ontario and Michigan, with most of the angled fish originating from wild populations, not hatcheries (Johnson et al. 2010). Feral populations of Chinook salmon likely vary in their degree of wildness, with some populations heavily influenced by hatchery supplementation and others not. We were able to estimate timeframes for colonisation of tributaries in different

regions of Lake Huron using commercial fisheries bycatch data and have provided new information about the level of genetic divergence among feral populations.

Pinpointing the exact year Chinook salmon colonised tributaries of Lake Huron is challenging because of the lack of any long term, standardised, stream monitoring programme. Moreover, genetic divergence among populations was too low to reconstruct colonisation history with precision. Commercial bycatch data, however, provided useful information because the presence of fish in an area at least suggests the potential for access to tributaries. The spatial distribution of the catch shifted over time, suggesting different colonisation patterns among regions. High catch rates of Chinook salmon in the Main Basin in 1975–1979

suggest the potential for colonisation of Main Basin tributaries, including the Maitland River, Carp River, and Nunn's Creek, as early as 1975. In the southern Georgian Bay region, commercial catch data in combination with stream assessments suggest a slightly later colonisation date of 1980. Age 0 fish were found in the Sydenham and Pottawatomi Rivers in 1983 (Kerr & Perron 1986), and Chinook salmon bycatch was observed in nearby fisheries in the early 1980s. The commercial bycatch data also suggest the potential for colonisation of North Channel tributaries in the early 1980s. Chinook salmon are effective colonisers, and by 1995–1999, they had expanded their range into all areas of Lake Huron. With a mean generation time of 3.2 years, populations likely have been reproducing for a maximum of 10 generations in Main Basin tributaries and eight generations in southern Georgian Bay and North Channel tributaries.

We detected emerging population structure of Chinook salmon in Lake Huron. Based on multiple lines of evidence, the Maitland River population showed the greatest degree of genetic distinctness from the remaining populations. In this population, all pairwise F_{ST} and R_{ST} values were significant, and individuals were assigned to a separate genetic cluster in the Bayesian analysis. Thus, significant genetic differentiation has occurred in fewer than 10 generations in Lake Huron Chinook salmon. The Root River also showed evidence of genetic differentiation based on a large number of significant pairwise F_{ST} and R_{ST} values. Finally, the Nunn's Creek population had evidence of genetic differentiation based on five significant pairwise F_{ST} and R_{ST} values and on their assignment to a separate cluster. Because Nunn's Creek is heavily supplemented by the State of Michigan using gametes from returns to the Swan River (where no natural reproduction occurs), its differentiation most likely reflects divergence of the Swan River stock from feral populations elsewhere in Lake Huron.

Founder effects likely account for some of the genetic differentiation we observed among feral populations. In addition, because F_{ST} is approximately equal to $1/(4Nm + 1)$ (Wright 1969), significant genetic structuring could arise because of low effective population sizes (N) or low migration rates between populations (m). With the exception of the few populations receiving hatchery-reared fish from non-natal sources, we expect m in our study system to be similar to published values. Pacific salmon imprint to cues from natal or release sites, resulting in low rates of straying ($m = 0.01–0.04$) (Quinn 2005). Assuming $m = 0.01–0.04$, our largest observed value of F_{ST} (0.13) would be expected for $N = 42–167$. Thus for reasonable values of m , small effective populations can lead to significant genetic differentiation because

of genetic drift. Moreover, compared to larger populations, smaller populations allow F_{ST} to increase from zero to $1/(4Nm + 1)$ in fewer generations (Allendorf & Phelps 1981). Unfortunately, data on effective population sizes are not available for our study system for comparison with these estimated values of N . We know the Nottawasaga River likely has the largest feral Chinook salmon population in Lake Huron, but little else. To summarise, founder effects followed by genetic drift in small populations likely explains the differentiation of the Maitland River and Root River populations.

The amount of genetic differentiation among the remaining collection sites in Lake Huron was low. Weeder et al. (2005) concluded that hatchery practices, in addition to recent colonisation, contributed to a lack of genetic differentiation in allozyme loci among feral Chinook salmon populations in Michigan, and the same argument could be made in our study. For example, the Sydenham River population, which is a significant source of gametes to hatcheries that release hatchery-reared juveniles into non-natal water bodies, was not genetically distinct from other populations. Not only does the Sydenham River contribute gametes to the LHFC hatchery, it contributes gametes to a hatchery in Sarnia, Ontario (south-eastern Main Basin), which releases age 0 juveniles at several sites in local streams and along the lakeshore. Adults returning to these release locations would not find spawning sites (there are no known significant spawning sites south of the Maitland River) and would likely stray elsewhere. Recent colonisation and continued hatchery supplementation may also explain the absence of any signal of genetic isolation by distance in our study.

In situations involving the colonisation of new habitats by invasive or transplanted populations, trait evolution can happen very rapidly (Stockwell et al. 2003). Studies of *Oncorhynchus* spp. have provided fundamental insights into the dynamics of contemporary adaptation. Following colonisation, divergent habitat-mediated selection can be sufficiently strong to detect trait evolution and local adaptation in as few as 13–26 generations (Hendry et al. 2000; Kinnison et al. 2001; Quinn et al. 2001). Adaptation is likely in Great Lakes populations because imprinting naturally favours a higher degree of reproductive isolation and environmental selection pressures likely would be different from those experienced by the ancestral, Green River population. In particular, selection should be responsive to altered seasonal patterns of stream temperature and flow (Jonsson & Jonsson 2009). To test for local adaptation in Lake Huron, further study could focus on the characterisation of functional, non-neutral genetic markers such as MHC (major histocompatibility complex). Further study of local

Population differentiation in introduced salmon

adaptation could also focus on phenotype-habitat correlations, reciprocal transplant experiments or formal analyses of selection (Endler 1986; Reznick & Travis 1996; Schlüter 2000). Adaptation might happen first in large populations, such as the Nottawasaga River, that may be less subject to genetic drift. Adaptation might also happen quickly at range limits where environmental selection pressures would be most divergent within the Great Lakes.

Until recently, Chinook salmon fisheries in the Great Lakes were believed to be sustained by hatchery supplementation programmes. It is now apparent that wild reproduction is rampant, and many feral populations may be self-sustaining (Johnson et al. 2010). In our study, we detected emerging population genetic structure among feral populations in Lake Huron. Local adaptation of reproductively isolated populations should favour the persistence of Chinook salmon in Lake Huron, and even more so if stocking programmes cease. While the recreational fishery may benefit by the continued availability of a desirable species, there is also the potential for negative ecological effects on native species such as brook trout (*Salvelinus fontinalis* Mitchell) (Janetski et al. 2011) and lake trout (Negus 1995; Kitchell et al. 2000) and unknown effects on aquatic species at risk such as lake sturgeon (*Acipenser fulvescens* Rafinesque), northern brook lamprey (*Ichthyomyzon fosser* Reighard and Cummins) and redside dace (*Clinostomus elongatus* Kirkland).

Acknowledgements

We thank A. Moerke (Lake Superior State University Aquatic Research Laboratory), L. Hillyer and R. Keetch (OMNR), the Lake Huron Fishing Club, and the Sydenham Sportsmen's Association for providing Chinook salmon samples, A. Cottrill (OMNR) for providing access to the commercial catch database, S. Y. Baek for help with laboratory work and D. Gonder (OMNR) and anonymous reviewers for constructive comments on the manuscript. Financial support was provided by the Great Lakes Fishery Commission (Fishery Research Program to YM and three others), University of Western Ontario (Academic Development Fund to YM and BN), and the Natural Sciences and Engineering Research Council of Canada (Discovery Grant to YM).

References

- Allendorf, F.W. & Phelps, S.R. 1981. Use of allele frequencies to describe population structure. Canadian Journal of Fisheries and Aquatic Sciences 38: 1507–1514.
- Balloux, F. & Lugon-Moulin, N. 2002. The estimation of population differentiation with microsatellite markers. Molecular Ecology 11: 155–165.
- Candy, J.R. & Beacham, T.D. 2000. Patterns of homing and straying in southern British Columbia coded-wire tagged Chinook salmon (*Oncorhynchus tshawytscha*) populations. Fisheries Research 47: 41–56.
- Coble, D.W., Bruesewitz, R.E., Fratt, T.W. & Scheirer, J.W. 1990. Lake trout, sea lampreys, and overfishing in the upper Great Lakes: a review and reanalysis. Transactions of the American Fisheries Society 119: 985–995.
- Corander, J., Waldmann, P., Marttinen, P. & Sillanpää, M.J. 2004. BAPS 2: enhanced possibilities for the analysis of genetic population structure. Bioinformatics 20: 2363–2369.
- Crawford, S.S. 2001. Salmonine introductions to the Laurentian Great Lakes: an historical review and evaluation of ecological effects. Ottawa, ON: NRC Press.
- Dittman, A.H. & Quinn, T.P. 1996. Homing in Pacific salmon: mechanisms and ecological basis. Journal of Experimental Biology 199: 83–91.
- Dobiesz, N.E., McLeish, D.A., Eshenroder, R.L., Bence, J.R., Mohr, L.C., Ebener, M.P., Nalepa, T.F., Woldt, A.P., Johnson, J.E., Argyle, R.L. & Makarewicz, J.C. 2005. Ecology of the Lake Huron fish community, 1970–1999. Canadian Journal of Fisheries and Aquatic Sciences 62: 1432–1451.
- Endler, J.A. 1986. Natural selection in the wild. Princeton, NJ: Princeton University Press.
- Eshenroder, R.L., Coble, D.W., Bruesewitz, R.E., Fratt, T.W. & Scheirer, J.W. 1992. Decline of lake trout in Lake Huron. Transactions of the American Fisheries Society 121: 548–554.
- Excoffier, L., Smouse, P.E. & Quattro, J.M. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. Genetics 131: 479–491.
- Gile, S.R. & Milne, S.W. 2006. Commercial catch sampling program: 2004 summary report. Upper Great Lakes Management Unit Lake Huron Report PS_LHA_CF04_001 (03–2006). Owen Sound, ON: Ontario Ministry of Natural Resources.
- Guo, W. & Thompson, E.A. 1992. A Monte Carlo method for combined segregation and linkage analysis. American Journal of Human Genetics 51: 1111–1126.
- Hansen, M.J. 1999. Lake trout in the Great Lakes: basinwide stock collapse and binational restoration. In: Taylor, W.W. & Ferreri, C.P., eds Great Lakes fisheries policy and management: a binational perspective. East Lansing, MI: Michigan State University Press, pp. 417–453.
- Hendry, A.P., Wenburg, J.K., Bentzen, P., Volk, E.C. & Quinn, T.P. 2000. Rapid evolution of reproductive isolation in the wild: evidence from introduced salmon. Science 290: 516–518.
- Hendry, A.P., Castric, V., Kinnison, M.T. & Quinn, T.P. 2004. The evolution of philopatry and dispersal: homing versus straying in salmonids. In: Hendry, A.P. & Stearns, S.C., eds Evolution illuminated: salmon and their relatives. Oxford: Oxford University Press, pp. 52–91.
- Janetski, D.J., Moerke, A.H., Chaloner, D.T. & Lamberti, G.A. 2011. Spawning salmon increase brook trout movements in a Lake Michigan tributary. Ecology of Freshwater Fish 20: 209–219.
- Johnson, J.E., DeWitt, S.P. & Gonder, D.J.A. 2010. Mass-marking reveals emerging self regulation of the Chinook salmon population in Lake Huron. North American Journal of Fisheries Management 30: 518–529.
- Jonsson, B. & Jonsson, N. 2009. A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo*

- salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology* 75: 2381–2447.
- Kalinowski, S.T. 2005. Do polymorphic loci require large sample sizes to estimate genetic distances? *Heredity* 94: 33–36.
- Kalinowski, S.T., Wagner, A.P. & Taper, M.L. 2006. ML-RELATE: a computer program for maximum likelihood estimation of relatedness and relationship. *Molecular Ecology Notes* 6: 576–579.
- Kerr, S.J. 1987. Chinook salmon, *Oncorhynchus tshawytscha*, smolting activity in five Lake Huron – Georgian Bay tributaries during the spring of 1986. Owen Sound, ON: Ontario Ministry of Natural Resources.
- Kerr, S.J. 2006. An historical review of fish culture, stocking and fish transfers in Ontario, 1865–2004. Peterborough, ON: Ontario Ministry of Natural Resources.
- Kerr, S.J. & Perron, D.A. 1986. An examination of Chinook salmon, *Oncorhynchus tshawytscha*, smolting activity in six selected Lake Huron-Georgian Bay tributaries. Owen Sound, ON: Ontario Ministry of Natural Resources.
- Kinnison, M.T., Unwin, M.J., Hendry, A.P. & Quinn, T.P. 2001. Migratory costs and the evolution of egg size and number in introduced and indigenous salmon populations. *Evolution* 55: 1656–1667.
- Kitchell, J.F., Cox, S.P., Harvey, C.J., Johnson, T.B., Mason, D.M., Schoen, K., Aydin, K., Bronte, C., Ebener, M., Hansen, M., Hoff, M., Schram, S., Schreiner, D. & Walters, C.J. 2000. Sustainability of the Lake Superior fish community: interactions in a food web context. *Ecosystems* 3: 545–560.
- Marklevitz, S.A.C., Fryer, B.J., Gonder, D., Yang, Z., Johnson, J., Moerke, A. & Morbey, Y.E. 2011. Use of otolith chemistry to discriminate juvenile Chinook salmon (*Oncorhynchus tshawytscha*) from different wild populations and hatcheries in Lake Huron. *Journal of Great Lakes Research*, DOI: 10.1016/j.jglr.2011.08.004.
- Michalakis, Y. & Excoffier, L. 1996. A generic estimation of population subdivision using distances between alleles with special reference for microsatellite loci. *Genetics* 142: 1061–1064.
- Miller, M.P. 1997. Tools for population genetic analysis (TFPGA) 1.3: a Windows program for the analysis of allozyme and molecular population genetic data. Available: <http://www.marksgeneticsoftware.net/>. Last accessed on 11 November 2011.
- Negus, M.T. 1995. Bioenergetics modeling as a salmonine management tool applied to Minnesota waters of Lake Superior. *North American Journal of Fisheries Management* 15: 60–78.
- Nei, M. 1987. Molecular evolutionary genetics. New York, NY: Columbia University Press.
- O'Connell, M., Danzmann, R.G., Cornuet, J.-M., Wright, J.M. & Ferguson, M.M. 1997. Differentiation of rainbow trout (*Oncorhynchus mykiss*) populations in Lake Ontario and the evaluation of the stepwise mutation and infinite allele mutation models using microsatellite variability. *Canadian Journal of Fisheries and Aquatic Sciences* 54: 1391–1399.
- van Oosterhout, C., Weetman, D. & Hutchinson, W.F. 2006. Estimation and adjustment of microsatellite null alleles in nonequilibrium populations. *Molecular Ecology Notes* 6: 255–256.
- O'Reilly, P.T., Herbinger, C. & Wright, J.M. 1998. Analysis of parentage determination in Atlantic salmon (*Salmo salar*) using microsatellites. *Animal Genetics* 29: 363–370.
- Peck, J.W., Jones, T.S., MacCallum, W.R. & Schram, S.T. 1999. Contribution of hatchery-reared fish to Chinook salmon populations and sport fisheries in Lake Superior. *North American Journal of Fisheries Management* 19: 155–164.
- Powell, M.J. & Miller, M. 1990. Shoal spawning by Chinook salmon in Lake Huron. *North American Journal of Fisheries Management* 10: 242–244.
- Quinn, T.P. 1993. A review of homing and straying of wild and hatchery-produced salmon. *Fisheries Research* 18: 29–44.
- Quinn, T.P. 2005. The behavior and ecology of Pacific salmon and trout. Bethesda, MD: American Fisheries Society. Seattle: University of Washington Press.
- Quinn, T.P., Kinnison, M.T. & Unwin, M.J. 2001. Evolution of Chinook salmon (*Oncorhynchus tshawytscha*) populations in New Zealand: pattern, rate, and process. *Genetica* 112–113: 493–513.
- Raymond, M. & Rousset, F. 1995. GENEPOL (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248–249.
- Reznick, D. & Travis, J. 1996. The empirical study of adaptation in natural populations. In: Rose, M.R. & Lauder, G.V., eds *Adaptation*. San Diego, CA: Academic Press, pp. 243–289.
- Schlüter, D. 2000. Ecology of adaptive radiation. Oxford: Oxford University Press.
- Schneider, S., Kueffer, J.J.-M., Roessli, D. & Excoffier, L. 1997. ARLEQUIN: an exploratory population genetics software environment. Computer software distributed by authors.
- Scribner, K.T., Gust, J.R. & Fields, R.L. 1996. Isolation and characterization of novel salmon microsatellite loci: cross-species amplification and population genetic applications. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 833–841.
- Smith, N.G., Sullivan, P.J. & Rudstam, L.G. 2006. Using otolith microstructure to determine natal origin of Lake Ontario Chinook salmon. *Transactions of the American Fisheries Society* 135: 908–914.
- Stockwell, C.A., Hendry, A.P. & Kinnison, M.T. 2003. Contemporary evolution meets conservation biology. *Trends in Ecology and Evolution* 18: 94–101.
- Weeder, J.A., Marshall, A.R. & Epifanio, J.M. 2005. An assessment of population genetic variation in Chinook salmon from seven Michigan rivers 30 years after introduction. *North American Journal of Fisheries Management* 25: 861–875.
- Weir, B.S. & Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution* 38: 1358–1370.
- Williamson, K.S., Cordes, J.F. & May, B. 2002. Characterization of microsatellite loci in Chinook salmon (*Oncorhynchus tshawytscha*) and cross-species amplification in other salmonids. *Molecular Ecology* 2: 17–19.
- Wright, S. 1969. Evolution and the genetics of populations. Vol. II. The theory of gene frequencies. Chicago, IL: University of Chicago Press.
- Young, F.W. 1996. Vista: the visual statistics system. Chapel Hill, NC: Thurstone Psychometric Laboratory Research Memorandum 94-1(c), University of North Carolina.