

ECOLOGY OF WESTERN NORTH AMERICA

EDITED BY

V.J. KRAJINA

VOL. 1.

DEPARTMENT OF BOTANY

UNIVERSITY of BRITISH COLUMBIA

VANCOUVER, B.C., CANADA, FEBRUARY 27, 1965

THE COASTAL WESTERN HEMLOCK ZONE ON THE SOUTH-WESTERN BRITISH COLUMBIA MAINLAND

Vegetation - Environment Patterns and Ecosystem Classification

László Orlóci

NATO Science Fellow, University College of North
Wales, Bangor, United Kingdom

The Climax Area Pattern

Since the time that Whitford and Craig (1918) described some forest zones ("climatic forest types") from British Columbia, several zonal classifications were proposed. These utilizing mainly climatic and vegetation criteria (Table I).

TABLE I. Zonal Classifications with Reference to the South Western British Columbia Mainland

Whitford and Craig 1918 (Climatic types)	Halliday 1937 (Sections)	Rowe 1959 (Sections)	Krajina 1959, 1964 (Biogeoclimatic units)
Douglas-fir -- Red cedar cedar -Hemlock	Madroño - Oak transition (C.1)	Straits of Georgia (C.1)	Coastal Douglas-fir Zone, Garry Oak (Dry) Subzone
	Southern Coast (C.2)	South Pacific Coast (C.2)	Coastal Douglas-fir Zone, Wet Subzone
			Coastal Western Hemlock Zone, Dry Subzone
Hemlock -Balsam			Coastal Western Hemlock Zone, Wet Subzone
The subalpine part of the "subalpine and muskeg type"		Coastal subalpine	Mountain Hemlock Zone
Hemlock - Sitka spruce	Part of the southern and central Coast (C.1, C.2) and northern Coast (C.4)	Part of the northern Pacific Coast (C.3), and Queen Charlotte Island (C.4)	Coastal Western Hemlock Zone, Sitka Spruce Subzone (edaphic; ocean salt spray)

The zonal classification concept has been broadened and deepened recently by Krajina (1959, 1964) and his students to include climate, soils, and vegetation as the three regular criteria for the recognition of biogeoclimatic regions, zones, and subzones.

The Coastal Western Hemlock Zone belongs to a perhumid mesothermal climate (Köppen Cfb) in which the zonal (or mesic) soils are strongly podzolized and western hemlock (*Tsuga heterophylla*) is the climatic climax tree species.

Two subzones are recognised within the Coastal Western Hemlock Zone: one dry and one wet. These sub-zones are distinct climax areas and constitute successive segments along a regional climatic gradient, separated on the basis of the climatic climax vegetation and zonal soils. Each subzone is characterized by specific tendencies in vegetation and soil development which converge to a hypothetical climatic climax (mesic) ecosystem.

Mesic soils are strongly podzolized in the Dry Subzone, characterized by the climatic climax *Tsuga heterophylla*-*Plagiothecium undulatum* association. Association is used in the sense of an abstract vegetation type. Mesic soils of the Wet Subzone are not only strongly podzolized but moderately gleyed. In this wettest belt of the region (Table II) the climatic climax is the *Tsuga heterophylla* -*Abies amabilis* - *Vaccinium alaskaense* - *Plagiothecium undulatum* association.

TABLE II. Altitudinal Occurrence of Maximum Precipitation as Specified. After Orloci 1964

Observation period	1961*	January 1961	Long-term averages	October 1961	July 1961
Altitude at which maximum precipitation occurred	2390 ft.	4830 ft.	2450 ft.	2660ft.	2060 ft

* A random year of the last 10.

The ecotone of the Dry and Wet Subzones occurs between the 100 and 110 in. isohyets, corresponding approximately to 1200 ft. to 1300 ft. elevation. The division between the Wet Subzone and the Mountain Hemlock Zone lies in the critical altitudinal belt of 3000 ft. to 3200 ft. on slopes, and slightly lower in the valleys. Above these elevations snow accumulation is considerable during the winter months, but below them snow accumulation becomes insignificant (cf. Peterson 1964, Brooke 1964). A total annual precipitation of 60-70 inches appears to be the critical limit which separates the Coastal Douglas-fir Zone from the Coastal Western Hemlock Zone. The differentiating characteristics of the subzones of the Coastal Western Hemlock Zone are shown in Table III.

TABLE III. Differentiating Characteristics of the Climax Sites within the Coastal Western Hemlock Zone and the Adjacent Zones on the Southwestern British Columbia Mainland

Climax area	Climatic type	Mean annual temperature (°F)	Annual total precipitation (in.)	Annual total snowfall (in.)	Altitude (ft.)	Zonal soil-forming processes	Climatic climax (mesic) association
Coastal Douglas-fir Zone	Humid (summer dry) mesothermal	over 48	below 65	below 30	0-500	Weak podzolization, very weak laterization, moderate melanization	<i>Pseudotsuga menziesii</i> - <i>Gaultheria shallon</i> - <i>Eurhynchium oregonum</i>
Dry Subzone of the Coastal Western Hemlock Zone	Perhumid mesothermal	46-48	65-110	30-80	0-1200	Strong podzolization	<i>Tsuga heterophylla</i> - <i>Plagiothecium undulatum</i>
Wet Subzone of the Coastal Western Hemlock Zone	Perhumid mesothermal	43-46	over 110	80-200	600-3000	Strong podzolization, moderate gleyization	<i>Tsuga heterophylla</i> - <i>Abies amabilis</i> - <i>Vaccinium alaskaense</i> - <i>Plagiothecium undulatum</i>
Lower Subzone of the Mountain Hemlock Zone	Perhumid (snowy) microthermal	below 43	over 75	over 200	over 3000	Strong podzolization, strong gleyization	<i>Tsuga mertensiana</i> - <i>Abies amabilis</i> - <i>Vaccinium alaskaense</i> - <i>Rhizidiopsis robusta</i>

TABLE IV. Ecosystem Types of the Rock Outcrop Land Type, Dry Subzone

Associations	Rhacomitrium canescens (Rh)	Cladonieta - Polytrichetum (Cl)	Danthonietum spicatae (Da)
Land form	Knoll	Table	Crevice
Relief shape	Convex	Convex	Concave
Slope gradient ^o	27 (10-40) 9*	16 (10-30) 8*	9 (3-21) 4*
Altitude ft.	620 (250-1000) 9*	610 (250-1000) 8*	740 (250-1200) 4*
Hygrotope	Very dry	Very dry	Very dry
Permeable mineral soil depth (cm.)	1.4 (0.5-3.0) 9*	2.8 (0.5-0) 8*	23.0 (13-33) 4*
pH			
L-F	4.7 (4.3-5.1) 3*	4.4 (4.3-4.5) 2*	4.5(4.1-4.7) 3*
A	4.4 (4.0-5.0) 3*	4.8 (4.7-4.9) 2*	4.5(4.1-4.6) 3*
B			4.8(4.2-5.5) 3*
C			
Humus form	Raw humus	Raw humus	Duff mull
Characteristic combination of species	Rhacomitrium canescens	Cladonia pacifica Polytrichum piliferum Polytrichum juniperinum Cladonia rangiferina Rhacomitrium canescens Dicranum scoparium Cladonia gracilis	Danthonia spicata Polytrichum piliferum Polytrichum juniperinum Rhacomitrium canescens Cladonia pacifica Cladonia rangiferina Cladonia gracilis
	Polytrichetum commune (Pc)	Gaultherietum shallonis (Ga)	Cladonieta - Pseudotsugetum (LG)
Land form	Crevice	Complex	Upper-slope
Relief shape	Concave	Convex	Convex
Slope gradient ^o	7 (5-10) 3*	10 (5-21) 8*	25 (8-40) 5*
Altitude ft.	730 (600-1000) 3*	690 (500-1000) 8*	990 (540-1400) 5*
Hygrotope	wet	Dry to moist	Dry
Permeable mineral soil depth (cm.)	21 (7-35) 3*	16 (6-30) 8*	9 (3-13) 5*
pH			
L-F	4.4	4.7 (4.4-4.9) 5*	3.8 (3.7-4.0) 5*
A	5.0	4.4 (4.1-4.9) 5*	3.6 (3.6-3.7) 3*
B	5.4	5.2 (4.3-5.9) 5*	3.6 (3.6-3.7) 3*
C			
Humus form	Duff mull	Raw humus	Raw humus
Site index ¹⁾	Douglas-fir w. hemlock w. redcedar		69 (60-78) 5* 66 (64-67) 5* 56 (54-69) 5*
Characteristic combination of species	Polytrichum commune	Gaultheria shallon Danthonia spicata Polytrichum commune Pleurozium schreberi Rhacomitrium canescens Hylocomium splendens Eurhynchium oregonum Cladonia pacifica Cladonia rangiferina Polytrichum piliferum Polytrichum juniperinum	Pseudotsuga menziesii Gaultheria shallon Eurhynchium oregonum Hylocomium splendens Cladonia pacifica Cladonia rangiferina Rhacomitrium canescens Polytrichum piliferum Pleurozium schreberi

* The number of measurements.

¹⁾ Site index in ft. per 100 years in all tables.

The climax area pattern shifts in space as a consequence of regional climate change. Hansen (1947) has described two major changes in the climate which occurred in the region since the last glaciation. A warm climate prevailed 8000 to 4000 years ago in which the Coastal Douglas-fir Zone reached its maximum distribution. A return to a cooler climate forced the migration of the Coastal Western Hemlock Zone and the Mountain Hemlock Zone to their present positions.

Ecosystem Classification

It is appropriate to separate land types from within each subzone on the basis of the origin and origin of the parent materials as the second step in the stratification. These land types are natural units and they constitute distinct segments of the vegetation and ecotope.

Ecosystem types¹ of the Rock Outcrop Land Type, Dry Subzone

The rock outcrop land type includes localities which lack continuous soil cover, and in which the ground surface is a complex of projecting peaks, knolls and crevices. The patterns of these different habitats have distinct plant associations. The peaks are characterized by a loosely integrated association of crustose and foliose lichens. The knolls and the side slopes of the peaks are colonized by a bryophyte association in which *Rhacomitrium canescens* dominates. The community of fruticose lichens and some bryophytes is transitional between the *Rhacomitrium canescens* and the *Gaultheria shallon* associations. The dry crevices are occupied by *Danthonia spicata*, different lichens, and bryophytes. *Polytrichum commune* is typical in the wet crevices.

The *Gaultheria shallon* association occurs on thick raw humus along the forest edge. The side slopes and depressions where soil development is the most advanced are occupied by a lithosolic forest. This is the only forest association in the Coastal Western Hemlock Zone in which Douglas-fir (*Pseudotsuga menziesii*) can be self-perpetuating without fire or logging. The ecosystem types of the rock outcrop land type are described in Table IV.

Ecosystem types of the Glacial Till Land Type

The glacial till land type includes all localities of glacial deposits except those of the swamps. Glacial till originated during the Seymour, Semiamu, Vashon, and Capilano glaciations (Armstrong 1956, 1957). The Seymour sediments are more than 30,000 years old. The Capilano sediments are approximately 11,000 years old.

Two major environmental gradients were identified through ordination techniques within the glacial till land type as underlying causes of variation in the vegetation: soil-moisture gradient, local climatic gradient.

Three distinct associations occur along the soil-moisture gradient of the Dry Subzone: *Gaultheria* of xeric habitats, *Plagiothecium* of mesic habitats, and *Polystichum* of hygric habitats. The occurrence or absence of *Mahonia nervosa* signifies the warm or cool exposures respectively within the *Gaultheria* and the *Plagiothecium* associations. Utilizing the occurrence or absence of *Mahonia nervosa*, the *Gaultheria* and the *Plagiothecium* associations are subdivided into the *Gaultheria*-*Mahonia*, orthic *Gaultheria*, *Plagiothecium*-*Mahonia*, and the orthic *Plagiothecium* ecosystem types. The orthic and *Mahonia* variations constitute successive segments along a local climatic gradient from cool to warm aspects. The *Polystichum* association is divided into two ecosystem types: the orthic *Polystichum* with melanized soils, and many nitrophilous species and the degraded *Polystichum* on well podzolized soils, including an admixture of raw humus inhabiting plants. A brief description of these ecosystem types is given in Table V.

The xeric, mesic, and hygric segments of the soil moisture gradient within the Wet Subzone are characterized respectively by the *Vaccinium*-*Gaultheria*, the *Vaccinium*-*Plagiothecium*, and the *Blechnum* associations. The *Vaccinium*-*Gaultheria* association consists of two distinct ecosystem types: the orthic *Vaccinium*-*Gaultheria* on shallow glacial tills, and the lithosolic *Vaccinium*-*Gaultheria* on lithic (AeC) soils (Lesko 1961). The *Vaccinium*-*Plagiothecium* association is subdivided into an orthic and a hygric segment. The latter is transitional between the mesic and wet habitats. The influence of a local climatic gradient is apparent within the *Blechnum* association. The orthic *Blechnum* ecosystem type is characteristic for low elevations in the Wet Subzone. The *Blechnum*-*Streptopus* ecosystem type occurs on the higher elevations in the Wet Subzone and in the Mountain Hemlock Zone (Peterson 1964). The ecosystem types of the Wet Subzone are described in Table VI.

² Ecosystem type includes a multitude of individual ecosystems among which variations in floristic structure are independent of the environmental variations.

TABLE V. Ecosystem Types of the Glacial Drift Land Type, Dry Subzone

Associations	Gaultherieto - Pseudotsugum		Tusgetum heterophyllae	
	Orthic Gaultheria (OG)	Gaultheria-Mahonia (G-M)	Orthic Plagiothecium (OP1)	Plagiothecium-Mahonia (P1-M)
Ecosystem type	Upper-slope Convex to straight	Upper-slope Convex to straight	Middle-slope Straight (to convex)	Middle-slope Straight *
Land form	12 (3-25) 8*	31 (18-40) 5*	6 (1-17) 14*	18 (1-31) 4*
Relief shape	1030 (750-1640) 8*	750 (450-900) 5*	780 (200-1050) 14*	540 (360-880) 4*
Slope gradient ^o	Moderately dry (to mesic)	Moderately dry (to mesic)	Mesic (to mod. dry or moist)	Mesic (to moderately dry)
Altitude ft.	40 (25-60) 4*	60 (40-80) 2*	31 (1-60) 7*	2
Hygrotope	62 (40-85) 4*	98+	71+	125+
Stoniness (%)	Nil	Nil	Nil	Nil
Permeable mineral soil depth (cm.)	8 (3-18) 4*	7 (5-8) 2*	5 (1-13) 7*	5
Endo-humus depth (cm.)	4.0 (3.4-4.5) 4*	4.3	4.1 (3.4-4.5) 5*	4.0
Fluviated horizon depth (cm.)	4.0 (3.7-4.3) 4*	4.4	4.2 (3.8-4.7) 5*	4.1
pH	5.2 (5.0-5.5) 4*	5.7	5.4 (5.3-5.6) 5*	5.5
	5.6 (5.5-5.7) 4*	?	5.7	5.5
Humus form	Raw humus	Raw humus	Raw humus	Raw humus
Site index	Douglas-fir western hemlock western redcedar	93 (78-109) 5*	142 (114-157) 14*	138 (117-155) 4*
		86 (73-91) 5*	121 (83-143) 14*	109 (99-122) 4*
		73 (70-78) 5*	88 (70-132) 14*	84 (76-91) 4*
Characteristic combination of species	Tsuga heterophylla Pseudotsuga menziesii Gaultheria shallon Eurhynchium oreganum Hylocomium splendens Plagiothecium undulatum	Tsuga heterophylla Pseudotsuga menziesii Gaultheria shallon Mahonia nervosa Eurhynchium oreganum Hylocomium splendens Plagiothecium undulatum	Tsuga heterophylla Pseudotsuga menziesii Plagiothecium undulatum Rhytidiadelphus loreus Hylocomium splendens Eurhynchium oreganum	Tsuga heterophylla Pseudotsuga menziesii Mahonia nervosa Plagiothecium undulatum Hylocomium splendens Eurhynchium oreganum

* The number of measurements.

TABLE V. (continued)

Ecosystem type	Associations	
	Tsugetum heterophyllae Eurhynchium-Mahonia (E-M)	Polysticheto - Thujetum plicatae Orthic Polystichum (OPo)
Land form	Middle-slope	Lower-slope
Relief shape	Straight *	Concave *
Slope gradient ^b	22 (2-33) 4*	16 (3-40) 15*
Altitude ft.	820 (450-950) 4*	840 (460-1740) 15*
Hygrotopo	Mesic (to moist)	Moist (to wet)
Stoniness (%)	45 (40-50) 2*	37 (0-75) 10*
Permeable mineral soil depth (cm.)	45+	60+
Endo-humus depth (cm.)	Nil	Nil
Eluviated horizon depth (cm.)	8 (0-15) 2*	3 (0-15) 10*
pH	4.0 2*	4.3 (3.4-5.7) 7*
	4.0 2*	4.2 (3.5-5.0) 7*
	5.3 (5.0-5.6) 2*	5.3 (5.1-5.5) 7*
	?	5.7 (5.3-6.0) 7*
Humus form	Raw humus	Raw humus
Site index	131(121-141) 4*	163 (143-180) 15*
	103(95-118) 4*	132 (102-180) 15*
	98(90-108) 4*	118 (96-170) 15*
Characteristic combination of species	Tsuga heterophylla Pseudotsuga menziesii Mahonia nervosa Eurhynchium oregonum Hylocomium splendens Plagiothecium undulatum	Tsuga heterophylla Thuja plicata Pseudotsuga menziesii Acer circinatum Polystichum munitum Dryopteris astricta Hylocomium splendens Plagiothecium undulatum Eurhynchium oregonum Rhytidia delphus loreus

Note: Data were compiled from the field notes of Lesko (1961), Orlovi (1961), and Eis (1962). pH was measured by Lesko (1961).

* The number of measurements.

TABLE VI. Ecosystem Types of the Glacial Drift Land Type, Wet Subzone

Associations	Gaultherieto - Tsugetum heterophyllae		Abieteto - Tsugetum heterophyllae	
	Orthic Vaccinium - Gaultheria (CV-G)	Lithosolic Vaccinium - Gaultheria (LV-G)	Orthic Vaccinium - Plagiothecium (CV-P1)	Hygic Vaccinium - Plagiothecium (HV-P1)
Ecosystem type	Upper-slope Convex	Upper-slope of hilltop Convex	Middle-slope Straight	Middle-slope Straight to slightly concave
Land form				
Relief shape				
Slope gradient ^o	11 (8-16) 3*	7 (0-10) 5*	16 (5-30) 5*	10 (0-18) 14*
Altitude (ft.)	1730 (1450-2000) 3*	1510 (590-1800) 5*	1680 (690-2330) 5*	1100 (550-2100) 14*
Hygrotope	Dry	Dry	Mesic	Moist
Stoniness (%)	7 (5-10) 3*	2 (0-10) 5*	55 (40-80) 5*	16 (0-50) 14*
Permeable mineral soil depth (cm.)	18 (15-21) 3*	10 (4-15) 5*	127 (90-145) 5*	40+
Endo-humus depth (cm.)	Nil	Nil	Nil	Nil
Eluviated horizon depth (cm.)	9 (7-10) 3*	10 (4-15) 5*	3 (2-5) 5*	2 (0-8) 14*
pH	3.4 (3.3-3.5) 2*	3.7 (3.5-4.0) 2*	3.8 (3.5-4.1) 5*	3.6 (3.3-3.9) 11*
	3.5 (3.4-3.5) 2*	3.6 (3.4-3.7) 2*	3.9 (3.8-4.1) 5*	3.8 (3.6-4.6) 11*
	4.3 (4.0-4.6) 2*	Nil	5.2 (5.0-5.7) 5*	5.2 (4.7-5.4) 11*
	?	?	Raw humus	5.4 (4.9-5.7) 11*
Humus form	Raw humus	Raw humus	Raw humus	Raw humus
Site index	75 (70-79) 3*	69 (60-80) 5*	123	127 (118-135) 14*
	62 (53-69) 3*	56 (39-75) 5*	104 (91-116) 5*	106 (88-134) 14*
	65 (48-84) 3*	53 (38-81) 5*	100 (85-113) 5*	108 (85-129) 14*
	59	40	94 (78-117) 5*	85 (73-129) 14*
	44	67	Nil	Nil
Characteristic combination of species	Tsuga heterophylla Thuja plicata Pseudotsuga menziesii Pinus monticola Chamaecyparis nootkatensis Vaccinium alaskaense Gaultheria shallon Rhytidopsis robusta Hylocomium splendens Plagiothecium undulatum	Tsuga heterophylla Thuja plicata Pseudotsuga menziesii Pinus monticola Chamaecyparis nootkatensis Vaccinium alaskaense Gaultheria shallon Rhytidopsis robusta Hylocomium splendens Plagiothecium undulatum	Tsuga heterophylla Thuja plicata Abies amabilis Vaccinium alaskaense Blechnum spicant Clintonia uniflora Plagiothecium undulatum Rhytidadelphus foreus Rhytidopsis robusta	Tsuga heterophylla Thuja plicata Abies amabilis Vaccinium alaskaense (Acer circinatum) Blechnum spicant Tiarella trifoliata Dryopteris austriaca (Athyrum filix-femina) (Streptopus amplexifolius) Plagiothecium undulatum Rhytidadelphus foreus

* The number of measurements.

TABLE VI. (continued)

Associations		Blechneto - Tsugetum heterophyllae	
Ecosystem type		Orthic Blechnum (OB)	Blechnum - Streptopus (B-S)
Land form		Lower-slope	Lower-slope
Relief shape		Concave	Concave
Slope gradient ^o		7 (2-10) 13*	8 (5-14) 5*
Altitude (ft.)		1370 (570-2730) 13*	2810 (2650-2950) 5*
Hygrotope		Wet	Wet
Stoniness (%)		29 (0-60) 12*	34 (25-50) 4*
Permeable mineral soil depth		35+	18+
Endo-humus depth (cm.)		1 (0-15) 12*	Nil
Eluviated horizon depth (cm.)		5 (0-13) 12*	6 (0-13) 4*
pH	L-F	3.8 (3.2-4.4) 6*	3.2
	A	3.8 (3.5-4.1) 6*	?
	B	4.8 (4.3-5.1) 6*	4.2
	C	5.2	5.7
Humus form		Raw humus	Greasy raw humus
Site index	Douglas-fir	125 (96-149) 13*	?
	western hemlock	118 (96-136) 13*	108 (97-123) 5*
	western redcedar	107 (85-149) 13*	112
	amabilis fir	115 (77-141) 13*	109 (92-122) 5*
	yellow cedar	Nil	Nil
Characteristic combination of species		<i>Tsuga heterophylla</i> <i>Thuja plicata</i> <i>Abies amabilis</i> <i>Vaccinium ovalifolium</i> <i>Vaccinium alaskaense</i> <i>Rubus spectabilis</i> <i>Sambucus pubens</i> (<i>Oplopanax horridus</i>) <i>Blechnum spicant</i> <i>Tiarella trifoliata</i> <i>Athyrium filix-femina</i> <i>Streptopus amplexifolius</i> <i>Dryopteris austriaca</i> (<i>Lysichitum americanum</i>) (<i>Polystichum munitum</i>) <i>Rhytidiadelphus loreus</i> <i>Hylocomium splendens</i> (<i>Pellia</i> sp.) (<i>Conocephalum conicum</i>) (<i>Eurhynchium stokesii</i>)	<i>Tsuga heterophylla</i> <i>Thuja plicata</i> <i>Abies amabilis</i> <i>Vaccinium ovalifolium</i> <i>Vaccinium alaskaense</i> <i>Rubus spectabilis</i> <i>Sambucus pubens</i> (<i>Oplopanax horridus</i>) <i>Blechnum spicant</i> <i>Tiarella trifoliata</i> <i>Streptopus amplexifolius</i> <i>Athyrium filix-femina</i> <i>Rubus pedatus</i> <i>Streptopus roseus</i> <i>Dryopteris austriaca</i> <i>Cornus canadensis</i> (<i>Lysichitum americanum</i>) <i>Rhytidiadelphus loreus</i> <i>Rhytidopsis robusta</i> <i>Plagiothecium undulatum</i>

Note: Data were compiled from the field notes of Lesko (1961), Orloci (1961), and Eis (1962).

* The number of measurements.

Ecosystem types of the Spring-Water Swamp Land Type

After water infiltrates the soils on the slope, it begins to move laterally under gravitational pressure over the impervious layer. If the impervious layer outcrops on the slope, then the overlying water comes to the surface at this point, saturates the soils downslope, and a spring-water swamp is formed. Two pedogenically distinct types of spring-water swamps are encountered in the Coastal Western Hemlock Zone: one on waterlogged mineral soils, and the other on waterlogged woody peat. Swamps on mineral soils are confined mainly to slopes, while woody peat occur in the depressions. The ecosystem types of the spring-water swamp land type are described in Table VII. It should be mentioned that the *Vaccinium*-*Lysichitum* type is characteristic for localities with intensive water movement. The *Lysichitum*-*Coptis* type is confined to the margins of Swagnum bogs with stagnant water, but exposed to runoff water or seepage from the adjacent slopes.

Ecosystem type of the Ravine Alluvium Land Type

In the glossary of geology and related sciences (American Geological Institute 1962), a ravine is defined as a "depression worn out by running water, larger than a gully and smaller than a valley". This term is applied in the present work to narrow valley-like depressions which were cut by running water in the glacial till, and

TABLE VII. Ecosystem Types of the Spring-water Swamp and Ravine Alluvium Land Type

Associations	Lysichiteto - Thujetum plicatae		Copteto - Thujetum plicatae		Oplopanaceto - Thujetum plicatae	
	Vaccinium-Lysichitum (V-Ly)		Lysichitum-Coptis (Ly-C)		Oplopanax-Adiantum (O-A)	
Land form	Lower-slope and outwash terrace		Depression		Ravine	
Relief shape	Concave		Straight		Concave	
Slope gradient ^o	2 (0-10) 10*		0		6 (4-10) 5*	
Altitude ft.	880 (700-160) 10*		820 (700-1000) 3*		906 (480-1800) 5*	
Hygrotopo	swampy		swampy		swampy	
Ground water table	Near surface, moving		Near surface, stagnant.		Near surface, intermittent overflow	
pH	4.3 (3.0-4.9) 7*				5.5	
	5.3 (4.6-6.1) 7*				5.8	
Humus form	Anmoor or peat anmoor		Peat anmoor		Anmoor or mull	
Site index	102 (88-130) 10*		98 (80-129) 3*		136 (130-150) 5*	
	101 (76-125) 10*		79 (70-98) 3*		137 (84-160) 5*	
					121 (100-44) 5*	
Characteristic combination of species	<p>Thuja plicata Picea sitchensis) Vaccinium alaskaense Rubus spectabilis Lysichitum americanum Blechnum spicant Dryopteris austriaca Tiarella trifoliata Athyrium filix-femina Conocephalum conicum Mnium punctatum Eurhynchium stokesii Pellia sp. Sphagnum squarrosum</p>		<p>Thuja plicata Pinus monticola Menziesia ferruginea Rubus spectabilis Vaccinium alaskaense Sambucus pubens Lysichitum americanum Coptis asplenifolia Coptis trifolia Mnium punctatum Sphagnum squarrosum</p>		<p>Thuja plicata Oplopanax horridus Ribes bracteosum Rubus spectabilis Vaccinium alaskaense Lysichitum americanum Adiantum pedatum Athyrium filix-femina Blechnum spicant Viola glabella Circaea alpina Gymnocarpium dryopteris Tiarella trifoliata Dryopteris austriaca Eurhynchium stokesii Conocephalum conicum Pellia sp. Sphagnum squarrosum Mnium punctatum Mnium memmesii</p>	

* The number of measurements.

which usually contain permanent or semi-permanent streamlets or creeks.

The *Thuja plicata* - *Oplopanax horridus* association which characterizes the ravine alluvium land type is remarkably uniform throughout both subzones, constituting a single ecosystem type. It extends into the Mountain hemlock zone where *Thuja plicata* is replaced by *Chamaecyparis nootkatensis*. A brief description is given in Table VII.

Ecosystem types of the Squamish Flood-Plain Land Type

Variations in floristic structure among the flood-plain ecosystem types are attributed mainly to differences in overflow and post-flood drainage. Overflow conditions are directly related to run-off induced fluctuation of the stream water level, and is therefore indirectly controlled by the climatic elements of the drainage area. Post-flood drainage reflects how rapidly the flood water is removed from an area when the water level of the stream starts to fall. In this connection, the differences are most marked between the benches along the active channel and the scars, between soils which contain a gravel horizon close to the soil surface, and soils which do not contain such a horizon, and so forth.

The terrain of a flood-plain aggrades by deposition of sediments from intermittent overflow. As the surface grows higher, overflow becomes less frequent. A similar reduction of overflow frequencies can occur as a consequence of downcutting in the river bed.

The bench height gradient coincides with a definite sequence of plant communities. Variations can occur at any particular bench height as a consequence of differences in post-flood drainage and soil quality. The ecosystem types of the Squamish flood-plain land type are described in Table VI. Scars and oxbow lakes are not discussed.

Plant communities are most specific in their ecotopic requirements in all land types which were considered in this paper. This makes it possible to utilize the plant communities as bioindicators of the different ecotopes.

Development of Ecosystems

Development implies vegetational and environmental ecosystem changes which result in a successional (time) sequence of plant communities and ecotopes at any particular point of the landscape. Developmental trends in time can be anticipated on the basis of present plant community distribution in space along existing environmental (eco-topical) gradients. The successional sequence, so defined, necessarily pertains only to plant communities and ecotopes which presently exist in the general region. It should be mentioned that a successional sequence of plant communities and ecotopes does not necessarily coincide with their geographic sequence.

Each individual plant community is visualized as the product of development that can be produced in a particular habitat. Any further plant community development requires a habitat change. In this way, each individual community represents a habitat "climax" ("permanent community" sensu Braun-Blanquet 1932). A geographic pattern of the habitat climax communities ("climax pattern" sensu Whittaker 1953) appears as a cross-section across individual developmental lines, evolving at different points of the landscape.

Lesko (1961) has recognized within the Coastal Western Hemlock Zone distinct trends in soil development that converge to mesic podzols. These podzols have a thick raw humus surface horizon, a prominent eluviated horizon (Ae) and a cemented illuvial horizon rich in organic, clay colloids, and iron oxides (Bfhc). The lower part of the B horizon is gleyed in the Wet Subzone, but gleization is absent from these soils in the Dry Subzone.

The convergence in plant community development to a climatic climax state is the consequence of a trend in habitat change towards a mesic condition under the given climate. The concept of successional convergence implied in the present work can be understood at two levels of generalization: a universal establishment of a particular dominance type appropriate to the forest climate, and the establishment of a mesic state of environment and vegetation through a sequence in time of many different plant associations at a given point of the landscape. While a trend of successional convergence is recognized, it does not mean that development must go through the complete successional sequence in every locality of the landscape. In some localities, development is very slow and it would require substantial physiographic changes to move toward a mesic status, longer than the persistence of the given climate. In a mountainous terrain substantial physiographic change, however, may involve a period of time far exceeding that required for substantial soil or climatic changes. The climax pattern therefore, may change geographically in such a way, that the developmental lines evolving at the different points of the landscape may be re-oriented in the direction of new attractor conditions several times before a climatic climax state ecosystem would develop in certain localities. No distinction is made between climax and seral communities in a sense that one is more stable than the other. The climax community itself is a changing entity, and therefore in successional convergence, as Cowles (1901) pointed out, one variable approaches another variable rather than a constant.

The direction in plant community development depends on the nature of the environmental gradient produced in a locality by environmental changes in time. For example, a change to a warmer and drier climate could result in successions as indicated by the horizontal arrow from right to left in Table X. This succession could lead, for example, to the establishment of plant communities which characterize the Coastal Douglas-fir Zone in place of those which occur at present in the Coastal Western Hemlock Zone.

TABLE VIII. Ecosystem Types of the Squamish Flood-plain Land Type, Benches

Associations	Hygrohypneto - Scoulerietum (H-S)	Equisetum arvense (E)	Scoulerieta - Salicetum (S-S)	Saliceto - Populetum (S-P)
Bench height (ft.)	6.5 (5-8) 4*	12 (11-13) 3*	10	11.9 (10-13.5) 8*
Days of overflow per year	146 (102-190) 4*	10 (4-18) 3*	42	15 (3-42) 8*
Depth to first gravel horizon (cm.)	0	44 (33-60) 3*	0	32 (0-95) 8*
Stand age (years)	0	--	?	4 (2-7) 8*
Characteristic combination of species	Scouleria aquatica Hydrohypnum ochraceum	Equisetum arvense	Salix sitchensis Alnus rubra Populus trichocarpa Elymus glaucus Scouleria aquatica	Populus trichocarpa Salix sitchensis Alnus rubra Salix lasiandra Equisetum arvense
Associations	Lonicereta - Populetum Lonicera - Rubus type (L-R)	Oplopanaceto - Piceetum Ribes - Oplopanax type (R-O)	Symphoricarpeto - Piceetum Symphoricarpos type (S)	
Bench height (ft.)	15.5 (14-17.5) 10*	19.6 (18-21.5) 4*		19.2 (18-21) 5*
Days of overflow per year	2 (up to 3) 10*	less than 1		less than 1
Depth to first gravel horizon (cm.)	104 (50-120) 10*	125+		69 (60-85) 5*
Stand age (years)	19 (14-45) 10*	66 (25-89) 4*		40 (27-55) 5*
Characteristic combination of species	Populus trichocarpa Alnus rubra Picea sitchensis (Thuja plicata) Salix sitchensis Lonicera involucrata Rubus spectabilis Maianthemum dilatatum Osmorhiza chilensis Equisetum arvense	Picea sitchensis Populus trichocarpa (Alnus rubra) (Acer macrophyllum) Thuja plicata Oplopanax horridus Ribes bracteatum Athyrum filix-femina (Dryopteris filix-mas) (Polystichum andersonii) Polystichum munitum Maianthemum dilatatum Osmorhiza chilensis Tiarella trifoliata Dryopteris austriaca Disporum oregonum Mnium insigne	Picea sitchensis (Acer macrophyllum) Thuja plicata Populus trichocarpa Symphoricarpos rivularis Disporum oregonum Polystichum munitum Athyrum filix-femina (Dryopteris filix-mas) Dryopteris austriaca Osmorhiza chilensis Mnium insigne	

* The number of measurements.

Primary successions on Rock Outcrops (Dry Subzone)

Since the loosened particles are immediately removed by wind, water, and gravity the steep rock outcrop, little essential endogenic development can occur in their ecosystem, and, the initial colonising stages can be greatly prolonged. Shade, litter, and root action of the arboreal vegetation around the pioneer communities, however, can accelerate development in them far beyond their endogenic potentials.

Development binds the pioneer communities into a common complex with *Gaultheria shallon*. At this stage of, organic matter accumulates more rapidly than in any one of the preceding stages, and raw humus (or root mor) is formed. Accumulation of raw humus and the root action of *Gaultheria* cause the destruction of the pioneer community fragments, and promote uniformity.

The decomposition of the rock surface is accelerated under the *Gaultheria* association. The weathering products are well protected against erosion, they accumulate, become leached, and form a soil horizon which is ash-gray in colour. The soil profile, hence, consists of a thick root mor underlain by an eluvial (Ae) horizon and the parent rock. This soil type was described by Lesko (1961) as an Eluviated Lithosol.

Establishment of Douglas-fir (*Pseudotsuga menziesii*) leads to the development of a lithosolic forest. In this stage, the disintegration of the remnants of the pioneer lichen and bryophyte communities is nearly complete, owing to shading and root action. Roots concentrate in the crevices, thereby occupying the space with the extra soil which had accumulated there and which conditioned the floristic structure of the *Danthonia spicata* and the *Polytrichum commune* associations. The sequence of primary successions on rock outcrops is shown in Table IX.

Successions in some forest ecosystems

Glacial till soils are very unstable on steep slopes. They are easily removed by erosion when the continuity of the vegetation cover is interrupted, especially after logging and fires. The trend is the complete removal of soils from the steep and convex slopes, and a subsequent expansion of the lithosolic forest (LG) and other rock outcrop communities. Accumulation of the weathering products is very slow and the erosion potential in the perhumid climate is extreme. Materials that are removed from the upper slopes are deposited on the lower slopes or carried further by the run off. These deposits increase soil depth and alter the soil-moisture regime. In this way, aggradation is a significant source of development on the lower slopes.

Glacial drift is a relatively young and little altered material in the Coastal forests of British Columbia. Therefore, soils originating from glacial drift can change a great deal in a relatively short period of time. The perhumid meso-thermal climate supports raw humus accumulation, leaching, and subsequent ortstein development. The ortstein layer may develop at various depths in the soil profile. This inevitably reduces the effective soil depth and alters the soil-moisture regime. In some habitats, seepage may occur in the at the impervious layer and succession may proceed towards hygro- or hydrophytic communities. Ortstein development may also bring about a shift toward xeric conditions in those soils which are intensively drained and lack seepage.

Seepage will contribute to changes in the soil by causing sedimentation of fine materials in the soil profile which, in turn, reduce aeration and impede drainage. Seepage water, enriched by nutrients, affects the floristic composition of the plant communities and consequently may influence the humus quality. The nutrient richness can promote fast decomposition of the organic matter, thereby affecting soil and plant community development. Successions in the forest ecosystems of the Coastal Western Hemlock Zone, except those of the flood plains, are outlined in Table X.


Successions on the Squamish flood plain

While the meander cuts away at the outer bank where the turbulence of the flow is greatest, it deposits sediments along the inner bank where flow velocity is least. This causes migration of the meander and builds up a new terrain that is open to plant colonization. Additional deposits of sediments will raise the surface of the terrain above a critical level where plant colonization can begin.


Different successional trends (see Table XI) evolve on the different terra nova. *Equisetum arvense* is the first species on loose sand. By its long rhizomes, *Equisetum arvense* efficiently stabilizes sand and promotes aggradation. *Scouleria aquatica* and *Hygrohypnum ochraceum* form an initial community on stabilized boulders. *Scouleria* and *Hygrohypnum* are attached firmly to the boulders and resist strong current and long summer overflow. Accumulation of a tiny horizon of silt from the withdrawing flood on gravel benches may provide a suitable seed-bed for the establishment of *Salix sitchensis*, *S. lasiandra*, *S. scouleriana*, *Alnus rubra*, and *Populus trichocarpa*. Successful establishment of the trees, however, requires the coincidence of several factors. The bench should be slightly higher than the average summer water level, the flood should have retreated leaving a moist nutritive surface soil behind at the time of seed dissemination, and germination should be followed by a long flood-free period for successful development of the seedlings.

A vegetated surface will aggrade faster than one without vegetation. This will initiate a process of differentiation of levels on a flood plain. Topographic forms which were produced by an even aggradation under stands of vegetation may be called benches and they can be classified by their relative height above the zero water mark, a water level which coincides with zero of the water gauge. The extent of a uniform bench usually coincides with a uniform even-aged stand of vegetation in primary successions under which deposits were laid down by intermittent overflow at a rate and extent influenced by the vegetation itself. Within these limits, soils, overflow conditions, and potential productivity are very similar. Different bench levels reflect variations in the time of establishment of the initial stand. The age of a primary stand is usually younger on a low bench than the age of a primary stand on a high bench.

TABLE IX. Primary Successions on Rock Outcrops



X. Successions in the Forest Ecosystems of the Coastal Western Hemlock Zone and the Adjacent Climax Areas




* After Krajina and Spilsbury (1953).

** After Peterson (1964) and Brooke (1964)

Note: Transitional and high moors are not considered.

TABLE XI. Primary Successions on the Squamish Flood-plain, Benches*



* Oxbow lakes are not discussed in this paper.

It is possible to distinguish between benches from the terraces in spite of their similarity in form. Hefley (1937; correctly pointed out that benches (or levels) are aggradation forms within a flood plain and they are attributable to the action of overflow water, wind, and vegetation. Terraces, on the other hand, mark a former valley floor level as stated by Thornbury (1958), and are largely the products of stream erosion rather than deposition (Gilbert 1877).

At the upper limit of the *Scouleria* - *Hydrohypnum* bench, the spaces among boulders are filled with sand and the establishment of willows, *Alnus rubra*, and *Populus trichocarpa* may take place. The boulders become fully or partially covered by sand while the bryophytes disappear gradually as they become buried. The establishment of *Elymus glaucus*, *Lonicera involucrata*, and *Rubus spectabilis* increases. At this stage the stand is 20 to 30 years old, *Populus trichocarpa* dominates in the crown canopy, and *Picea sitchensis* is abundant in the shrub layer. The soil is from two to three feet deep and consists of sand underlain by gravel or stones.

On the *Equisetum arvense* bench, development leads also to a *Lonicera* - *Rubus* community, in which the soil is similar to that just described except that boulders are not present in the subsoil and the sand is somewhat deeper. Development on the gravel benches converges also to a *Lonicera* - *Rubus* community.

The *Lonicera* - *Rubus* bench is frequently flooded but the deposited sediments are finer than before. Eventually a loamy sand surface horizon accumulates and the proportion of *Lonicera involucrata* decreases. The establishment of *Symphoricarpos rivularis* on this bench level indicates that a gravel horizon is close to the soil surface and follows a recent shift in the position of the river bed. *Oplopanax horridum* and *Ribes bracteosum* are specific to deep loam soils. In both the *Symphoricarpos* and the *Ribes* - *Oplopanax* community, *Picea sitchensis* and *Populus trichocarpa* dominate at the beginning, and subsequently *Picea sitchensis* alone. On this level, overflow is extremely rare and vegetation is conditioned by underground movement of water which varies with the water level of the stream.

Development on the *Symphoricarpos* and the *Ribes* - *Oplopanax* bench is very slow and if the habitat is no longer influenced by flood-water but seepage is present, the stand will remain rich in deciduous trees and herbs. These conditions promote an efficient decomposition of the organic matter and melanization. If no seepage is present in the soil, the stand will be dominated by conifers and raw humus accumulates. This leads to podzolization of the soil. Melanization and seepage are associated with the abundant occurrence of *Polystichum munitum*, and the podzols with raw humus inhabiting plants.

Ecosystem patterns of a flood plain are in a continuous flux. As the meanders migrate from one side of the valley to the other, benches and plant communities are destroyed along the outside bank. Development of an alternate channel can initiate substantial changes in the floristic structure of the plant communities through changes in the patterns of overflow and post-flood drainage. The distance of flow is always shortened by the alternate channel; this increases the river gradient. A new equilibrium is achieved by downcutting in the river bed.

If the break-through occurs across a medium bench, a unique situation can develop. Overflow duration is drastically reduced on the benches due to downcutting which results in a relative increase of bench height. Soil depth remains as before because bench level is increased by downcutting and not by sedimentation. The lesser vegetation will undergo rapid changes but the crown canopy will remain essentially unaltered for a long period of time depending on the stand age.

The overflow pattern in the unique situation just described resembles that of the *Ribes* - *Oplopanax* type. The soil profile which contains a gravel horizon close to the soil surface is similar to that of the *Lonicera* - *Rubus* type. The rate of sedimentation is low, and therefore no significant increase in soil depth can be anticipated. The crown canopy consists of deciduous trees in the early developmental stages (as in the *Lonicera* - *Rubus* type) but these give way later to the conifers (as in the *Ribes* - *Oplopanax* type). In the lesser vegetation, *Symphoricarpos rivularis* is the most abundant species.

The *Symphoricarpos* type always indicates a gravel horizon in the soil profile close to the soil surface, but this horizon does not necessarily indicate an earlier downcutting in the river bed. The torrential tributary streams can dump gravel even on the high benches of the main stream.

REFERENCES

- Armstrong, J. E. 1956. Surficial geology of Vancouver area, B. C. Geol. Surv. Can. Paper 55-40.
- _____. 1957. Surficial geology of New Westminster Map-area, B. C. Geol. Surv. Can. Paper 57-5.
- Braun-Blanquet, J. 1932. Plant sociology, the study of plant communities. (Trans. by G. D. Fuller and H. S. Conard.) New York McGraw-Hill.
- Brooke, R. C. 1964. Personal communications.
- Cowles, H. C. 1901. The physiographic ecology of Chicago and vicinity; a study of the origin, development, and classification of plant societies. Bot. Gaz. 31:108, 145-182.
- Eis, S. 1962. Statistical analysis of tree growth and some environmental factors of plant communities. Ph. D. Thesis, Dept. Biol. and Bot., Univ. of B. C.
- Gilbert, G. K. 1877. Report on the geology of the Henry Mountains. U. S. Geol. Geogr. Surv. of the Rocky Mountain Region (Powell) Washington: 120-127.

- Halliday, W. E. D. 1937. A forest classification of Canada. For. Serv. Bul. 89. Can. Dept. Mines and Res. Ottawa.
- Hansen, H. P. 1947. Climate versus fire and soil as factors in post-glacial forest succession in the Puget lowland of Washington. Am. Jour. Sci. 245: 265-286.
- Hefley, H. M. 1937. Ecological studies of the South Canadian river flood-plain in Cleveland County, Oklahoma. Ecol. Monogr. 7:345-402.
- Krajina, V. J. 1959. Bioclimatic zones in British Columbia. Univ. of B. C., Bot. Ser. No. 1, Vancouver, B. C.
- _____. 1964. Ecology of the forests of the Pacific Northwest. 1963 Prog. Report, Nat. Res. Coun. Grant T-92, 71-87.
- _____ and L. Orłóci. 1963. Biogeoclimatic forest zones on the lower mainland of British Columbia. 1962 Prog. Report, Nat. Res. Coun. Grant T-92, 18-24.
- _____, and R. H. Spilsbury. 1953. Forest associations on the east coast of Vancouver Island. Forestry Handbook for British Columbia, 1st. ed. (1953). Forest Club, University of British Columbia.
- Lesko, G. 1961. Ecological study of soils in the Coastal Western Hemlock Zone. M. Sc. Thesis, Dept. Biol. and Bot., Univ. of B. C.
- Orłóci, L. 1964. Vegetational and environmental variations in the ecosystems of the Coastal Western Hemlock Zone. Ph.D. Thesis, Dept. Biol. Bot., Univ. of B. C.
- Peterson, E. B. 1964. Plant associations in the Subalpine Mountain Hemlock Zone in southern British Columbia. Ph.D. Thesis, Dept. Biol. and Bot., Univ. of B. C.
- Rowe, J. S. 1959. Forest regions of Canada. For. Br., Bul. 123. Can. Dept. Nor. Aff. and Nat. Res. Ottawa. Thornbury, W.D. 1954, 1958, 1962. Principles of geomorphology. John Wiley & Sons, Inc. New York and London. Whitford, H. N. and R. D. Craig. 1918. Forests of British Columbia. Can. Com. Cons. Ottawa.
- Whittaker, R. H. 1953. A consideration of climax theory: the climax as a population and pattern. Ecol. Monogr. 23:41-78.

