

# Basic Model Theory

John L. Bell

## 1. Structures and First-Order Languages

A *structure* is a triple

$$\mathfrak{A} = (A, \{R_i: i \in I\}, \{e_j: j \in J\}),$$

where  $A$ , the *domain* or *universe* of  $\mathfrak{A}$ , is a *nonempty* set,  $\{R_i: i \in I\}$  is an indexed family of relations on  $A$  and  $\{e_j: j \in J\}$  is an indexed set of elements —the *designated elements* of  $A$ . For each  $i \in I$  there is then a natural number  $\lambda(i)$  —the *degree* of  $R_i$  —such that  $R_i$  is a  $\lambda(i)$ -place relation on  $A$ , i.e.,  $R_i \subseteq A^{\lambda(i)}$ . This  $\lambda$  may be regarded as a function from  $I$  to the set  $\omega$  of natural numbers; the pair  $(\lambda, J)$  is called the *type* of  $\mathfrak{A}$ . Structures of the same type are said to be *similar*.

Note that since an  $n$ -place operation  $f: A^n \rightarrow A$  can be regarded as an  $(n+1)$ -place relation on  $A$ , algebraic structures containing operations such as groups, rings, vector spaces, etc. may be construed as structures in the above sense.

The *cardinality*  $\|\mathfrak{A}\|$  of a structure  $\mathfrak{A}$  is defined to be the cardinality  $|A|$  of its domain  $A$ .

The *first-order language*  $\mathcal{L}$  of type  $(\lambda, J)$  has the following categories of *basic symbols*:

- (i) *individual variables*: a denumerable sequence  $v_0, v_1, \dots$ ;
- (ii) *predicate symbols*: for each  $i \in I$ , a predicate symbol  $P_i$  of degree  $\lambda(i)$ ;
- (iii) *individual constants*: for each  $j \in J$  an individual constant  $c_j$ ;
- (iv) *equality symbol*: the symbol  $=$ ;
- (v) *logical operators*:  $\neg$  (negation),  $\wedge$  (conjunction);
- (vi) *existential quantifier symbol*:  $\exists$  ("there exists");
- (vii) *punctuation symbols*: e.g.  $( )$ ,  $[ ]$ .

Predicate and constant symbols are often called *extralogical* symbols; variables and constants are collectively known as *terms*: we shall use symbols  $t, u$ , possibly with subscripts, to denote arbitrary terms.

*Atomic formulas* of  $\mathcal{L}$  are finite strings of basic symbols of either of the forms  $P_i t_1 \dots t_{\lambda(i)}$  or  $t = u$ , where  $t_1, \dots, t_{\lambda(i)}, t, u$  are terms. *Formulas* of  $\mathcal{L}$  (or  $\mathcal{L}$ -*formulas*) are finite strings of basic symbols defined in the following recursive manner:

- (a) any atomic formula is a formula;
- (b) if  $\varphi, \psi$  are formulas, so also are  $\neg\varphi, \varphi \wedge \psi$ , and  $\exists x\varphi$ , where  $x$  is any variable  $v_n$ ;
- (c) a finite string of symbols is a formula exactly when it follows from finitely many applications of (a) and (b) that it is one.

We write  $Form(\mathcal{L})$  for the set of all formulas of  $\mathcal{L}$ . The *degree* (of complexity) of a formula is

defined to be the number of occurrences of logical operators and quantifiers in it.

The symbols  $\vee$  (disjunction),  $\rightarrow$  (implication) and  $\forall$  (universal quantifier) are introduced as *abbreviations*:

$$\begin{aligned} \varphi \vee \psi & \text{ for } \neg(\neg\varphi \wedge \neg\psi) \\ \varphi \rightarrow \psi & \text{ for } \neg\varphi \vee \psi \\ \varphi \leftrightarrow \psi & \text{ for } (\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi) \\ \forall x\varphi & \text{ for } \neg\exists x\neg\varphi. \end{aligned}$$

We also write  $\bigwedge_{i=1}^n \varphi_i$  for  $\varphi_1 \wedge \dots \wedge \varphi_n$ .

It will be assumed that the notions of *free* and *bound* occurrence of a variable in a formula are understood. We write  $\varphi(v_0, \dots, v_n)$  to indicate that the free variables of  $\varphi$  are among  $v_0, \dots, v_n$ . We also write  $\varphi(x/t)$ , or simply  $\varphi(t)$ , for the result of substituting  $t$  at each free occurrence of  $x$  in  $\varphi$ . More generally, we write  $\varphi(t_0, \dots, t_n)$  for the result of substituting  $t_i$  at each occurrence of  $v_i$ , for  $i = 0, \dots, n$ , in  $\varphi(v_0, \dots, v_n)$ . An  $\mathcal{L}$ -sentence is an  $\mathcal{L}$ -formula without free variables. We write  $Sent(\mathcal{L})$  for the set of all  $\mathcal{L}$ -sentences.

The *cardinality*  $\|\mathcal{L}\|$  of  $\mathcal{L}$  is defined to be the cardinality of its set of basic symbols.

**Lemma.**  $\|\mathcal{L}\| = |Form(\mathcal{L})|$ .

**Proof.** Let  $\|\mathcal{L}\| = \kappa$ . Since  $\kappa$  is infinite and each formula is a finite string of symbols,  $|Form(\mathcal{L})| \leq \kappa$ . The fact that  $\kappa$  is infinite also implies that either the set of terms or the set of predicate symbols of  $\mathcal{L}$  (or both) must have cardinality  $\kappa$ . In either case the set of atomic formulas of the form  $Pit\dots t$  has cardinality  $\kappa$ , so that  $|Form(\mathcal{L})| \geq \kappa$ . The Lemma follows. ■

For  $\Sigma \subseteq Sent(\mathcal{L})$  we define  $\mathcal{L}_\Sigma$  to be the language whose extralogical symbols are precisely those occurring in at least one sentence of  $\Sigma$ .

**Lemma.**  $\|\mathcal{L}_\Sigma\| = \max(\aleph_0, |\Sigma|)$ .

**Proof.** If  $\Sigma$  is finite, evidently  $\|\mathcal{L}_\Sigma\| = \aleph_0$ . Now suppose that  $|\Sigma| = \kappa \geq \aleph_0$ . We have  $|\Sigma| \leq |Form(\mathcal{L}_\Sigma)| = \|\mathcal{L}_\Sigma\|$  by the previous lemma. For each  $\sigma \in \Sigma$  let  $S(\sigma)$  be the set of ( $\mathcal{L}_\Sigma$ -) symbols occurring in  $\sigma$ : then  $S(\sigma)$  is finite. Also the set  $K$  of terms of  $\mathcal{L}_\Sigma$  is included in the union of the sets  $S(\sigma)$  for  $\sigma \in \Sigma$ , so that

$$|K| \leq |\bigcup\{S(\sigma) : \sigma \in \Sigma\}| \leq \sum_{\sigma \in \Sigma} |S(\sigma)| \leq |\Sigma| \cdot \aleph_0 = |\Sigma|.$$

Thus  $\|\mathcal{L}_\Sigma\| \leq |K| + \aleph_0 + \aleph_0 \leq |\Sigma|$ , and hence  $\|\mathcal{L}_\Sigma\| = |\Sigma|$  as required. ■

## 2. Satisfaction, validity, and models.

If  $\mathcal{L}$  is a first-order language, a structure having the same type as that of  $\mathcal{L}$  is called an  $\mathcal{L}$ -

structure. Let  $\mathfrak{A} = (A, \{R_i: i \in I\}, \{e_j: j \in J\})$  be an  $\mathcal{L}$ -structure, where  $\mathcal{L}$  has type  $(\lambda, J)$ , and let  $\mathbf{a} = (a_0, a_1, \dots)$  be a countable sequence of elements of  $A$  (such a sequence will be referred to henceforth as an  $A$ -sequence). For any predicate symbol or term of  $\mathcal{L}$ , we define its *interpretation under*  $(\mathfrak{A}, \mathbf{a})$  as follows:

$$P_i^{(\mathfrak{A}, \mathbf{a})} = R_i \quad c_j^{(\mathfrak{A}, \mathbf{a})} = e_j \quad v_n^{(\mathfrak{A}, \mathbf{a})} = a_n.$$

Since  $P_i^{(\mathfrak{A}, \mathbf{a})}$  and  $c_j^{(\mathfrak{A}, \mathbf{a})}$  depend only on  $\mathfrak{A}$ , we usually just write  $P_i^{\mathfrak{A}}$  and  $c_j^{\mathfrak{A}}$  for these and call them the *interpretations* of  $P_i$  and  $c_j$ , respectively, in  $\mathfrak{A}$ .

For  $n \in \omega$ ,  $b \in A$  we define

$$[n/b]\mathbf{a} = (a_0, a_1, \dots, a_{n-1}, b, a_{n+1}, \dots).$$

For  $\varphi \in \text{Form}(\mathcal{L})$  we define the relation  $\mathbf{a}$  satisfies  $\varphi$  in  $\mathfrak{A}$ , written

$$\mathfrak{A} \models_{\mathbf{a}} \varphi,$$

recursively on the degree of  $\varphi$  as follows:

1) for terms  $t, u$ ,

$$\mathfrak{A} \models_{\mathbf{a}} t = u \Leftrightarrow t^{(\mathfrak{A}, \mathbf{a})} = u^{(\mathfrak{A}, \mathbf{a})};$$

2) for terms  $t_1, \dots, t_{\lambda(i)}$ ,

$$\mathfrak{A} \models_{\mathbf{a}} P_i t_1 \dots t_{\lambda(i)} \Leftrightarrow R_i(t_1^{(\mathfrak{A}, \mathbf{a})}, \dots, t_{\lambda(i)}^{(\mathfrak{A}, \mathbf{a})});$$

3)  $\mathfrak{A} \models_{\mathbf{a}} \neg \varphi \Leftrightarrow \text{not } \mathfrak{A} \models_{\mathbf{a}} \varphi$ ;

4)  $\mathfrak{A} \models_{\mathbf{a}} \varphi \wedge \psi \Leftrightarrow \mathfrak{A} \models_{\mathbf{a}} \varphi$  and  $\mathfrak{A} \models_{\mathbf{a}} \psi$ ,

5)  $\mathfrak{A} \models_{\mathbf{a}} \exists v_n \varphi \Leftrightarrow$  for some  $b \in A$ ,  $\mathfrak{A} \models_{[n/b]\mathbf{a}} \varphi$ .

The following facts are then easily established:

(a)  $\mathfrak{A} \models_{\mathbf{a}} \forall v_n \varphi \Leftrightarrow$  for all  $b \in A$ ,  $\mathfrak{A} \models_{[n/b]\mathbf{a}} \varphi$ ;

(b) suppose that  $\mathbf{a}, \mathbf{b}$  are  $A$ -sequences such that  $a_n = b_n$  whenever  $v_n$  occurs free in  $\varphi$ .

Then

$$\mathfrak{A} \models_{\mathbf{a}} \varphi \Leftrightarrow \mathfrak{A} \models_{\mathbf{b}} \varphi,$$

In view of fact (b), the truth of  $\mathfrak{A} \models_{\mathbf{a}} \varphi$  depends only on the interpretations under  $(\mathfrak{A}, \mathbf{a})$  of the free variables of  $\varphi$ , that is, if these are among  $v_0, \dots, v_n$ , only on  $a_0, \dots, a_n$ . Accordingly, under these conditions we shall often write

$$\mathfrak{A} \models_{\mathbf{a}} \varphi[a_0, \dots, a_n] \quad \text{for } \mathfrak{A} \models_{\mathbf{a}} \varphi.$$

We say that a formula  $\varphi$  is *valid* in  $\mathfrak{A}$  if  $\mathfrak{A} \models_{\mathbf{a}} \varphi$  for every  $A$ -sequence  $\mathbf{a}$  and *satisfiable* in  $\mathfrak{A}$  if  $\mathfrak{A} \models_{\mathbf{a}} \varphi$  for some  $A$ -sequence  $\mathbf{a}$ . It follows from (b) above that a sentence  $\sigma$  is satisfiable in a given structure iff it is valid there. If  $\sigma$  is valid in  $\mathfrak{A}$  we write

$$\mathfrak{A} \models \sigma$$

and say that  $\mathfrak{A}$  is a *model* of  $\sigma$ , or that  $\sigma$  *holds* in  $\mathfrak{A}$ . If  $\Sigma \subseteq \text{Sent}(\mathcal{L})$ , we say that  $\mathfrak{A}$  is a *model* of  $\Sigma$ , and write

$$\mathfrak{A} \models \Sigma,$$

if  $\mathfrak{A}$  is a model of each member of  $\Sigma$ . If  $\varphi \in \text{Form}(\mathcal{L})$ , we say that  $\Sigma$  *logically entails*  $\varphi$ , and write

$$\Sigma \models \varphi,$$

if  $\varphi$  is valid in every model of  $\Sigma$ . In particular, we write

$$\models \varphi$$

for  $\emptyset \models \varphi$ ; a formula  $\varphi$  satisfying this condition is then valid in every ( $\mathcal{L}$ -) structure and is called *universally valid*.

Let  $\mathcal{L}^*$  be a language which is an *extension* of  $\mathcal{L}$ , i.e. obtained from  $\mathcal{L}$  by adding a set  $\{P_i: i \in I^*\}$  of new predicate symbols and a set  $\{c_j: j \in J^*\}$  of new constant symbols. Given an  $\mathcal{L}^*$ -structure

$$\mathfrak{A}^* = (A, \{R_i: i \in I \cup I^*\}, \{e_j: j \in J \cup J^*\}),$$

the  $\mathcal{L}$ -structure

$$\mathfrak{A} = (A, \{R_i: i \in I\}, \{e_j: j \in J\})$$

is called the  $\mathcal{L}$ -*reduction* of  $\mathfrak{A}^*$ . Analogously,  $\mathfrak{A}^*$  is called an  $\mathcal{L}^*$ -*expansion* of  $\mathfrak{A}$ . Notice that, while an  $\mathcal{L}^*$ -structure always has a unique  $\mathcal{L}$ -reduction, an  $\mathcal{L}$ -structure has in general more than one  $\mathcal{L}^*$ -expansion. We write  $\mathfrak{A}^*|_{\mathcal{L}}$  for the  $\mathcal{L}$ -reduction of  $\mathfrak{A}^*$ . It is important to keep in mind the fact that *expanding or reducing has no effect on the domain of a structure; these operations merely add or subtract relations and designated elements*.

The following lemmas are routine. The first is proved by a straightforward induction on the degree of complexity of formulas, the second follows from the definition of  $\models$ .

**Expansion lemma.** Let  $\Sigma \subseteq \text{Sent}(\mathcal{L})$ , let  $\mathcal{L}^*$  be any extension of  $\mathcal{L}$ , let  $\mathfrak{A}$  be any  $\mathcal{L}$ -structure, and let  $\mathfrak{A}^*$  be any  $\mathcal{L}^*$ -expansion of  $\mathfrak{A}$ . Then

$$\mathfrak{A} \models \Sigma \Leftrightarrow \mathfrak{A}^* \models \Sigma. \quad \blacksquare$$

**Constants lemma.** Let  $\mathfrak{A}$  be an  $\mathcal{L}$ -structure, let  $\varphi(v_0, \dots, v_n) \in \text{Form}(\mathcal{L})$ , and let  $c_0, \dots, c_n$  be constant symbols of  $\mathcal{L}$ . Then

$$\mathfrak{A} \models \varphi(c_0, \dots, c_n) \Leftrightarrow \mathfrak{A} \models \varphi[c_0^{\mathfrak{A}}, \dots, c_n^{\mathfrak{A}}]. \quad \blacksquare$$

### 3. Review of first-order predicate logic.

Let  $\mathcal{L}$  be a first-order language of type  $(\lambda, J)$ . We specify *axioms* and *rules of inference* for  $\mathcal{L}$  as follows. As *axioms* we take

- 1) all instances of propositional tautologies;
- 2) *equality axioms*:

$$t = t \quad t = u \rightarrow u = t \quad t = u \wedge u = v \rightarrow t = v \\ (t_1 = u_1 \wedge \dots \wedge t_{\lambda(i)} = u_{\lambda(i)}) \rightarrow [P_i t_1 \dots t_{\lambda(i)} \rightarrow P_i u_1 \dots u_{\lambda(i)}]$$

- 3) all formulas of the form

$$\forall x \varphi(x) \rightarrow \varphi(t) \quad \varphi(t) \rightarrow \exists x \varphi(x)$$

where, if  $t$  is a variable, it does not occur bound in  $\varphi$ .

The *rules of inference* of  $\mathcal{L}$  are:

1) *modus ponens*:

$$\frac{\varphi \quad \varphi \rightarrow \psi}{\psi}$$

2) *quantifier rules*: if  $x$  is not free in  $\varphi$ ,

$$\frac{\varphi \rightarrow \psi(x)}{\varphi \rightarrow \forall x\varphi(x)} \qquad \frac{\psi(x) \rightarrow \varphi}{\exists x\psi(x) \rightarrow \varphi}$$

A *proof* in  $\mathcal{L}$  of  $\varphi$  from a set  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is a finite sequence  $\psi_1, \dots, \psi_n$  of  $\mathcal{L}$ -formulas, with  $\psi_n = \varphi$ , each member of which is either an axiom, a member of  $\Sigma$ , or else follows from previous  $\psi_i$  by one of the rules of inference. We say that  $\varphi$  is *provable from*  $\Sigma$ , and write

$$\Sigma \vdash \varphi,$$

if there is a proof of  $\varphi$  from  $\Sigma$ .  $\Sigma$  is said to be *consistent* (in  $\mathcal{L}$ ) if for no  $\mathcal{L}$ -formula  $\varphi$  do we have  $\Sigma \vdash \varphi \wedge \neg\varphi$ . If  $\emptyset \vdash \varphi$ , we write  $\vdash \varphi$  and say that  $\varphi$  is a *theorem* of  $\mathcal{L}$ .

We now list a number of basic results concerning these notions. Throughout,  $\Sigma$  denotes an arbitrary set of  $\mathcal{L}$ -sentences.

**Quantifier lemma.** If  $x$  does not occur free in  $\varphi$ , then

$$\Sigma \vdash \exists x(\varphi \wedge \psi) \leftrightarrow (\varphi \wedge \exists x\psi) \quad \Sigma \vdash \exists x(\varphi \rightarrow \psi) \leftrightarrow (\varphi \rightarrow \exists x\psi). \quad \blacksquare$$

**Deduction theorem.** If  $\sigma \in \text{Sent}(\mathcal{L})$ , then for any formula  $\varphi$ ,

$$\Sigma \cup \{\sigma\} \vdash \varphi \leftrightarrow \Sigma \vdash \sigma \rightarrow \varphi. \quad \blacksquare$$

**Finiteness theorem.** If  $\Sigma \vdash \varphi$ , then  $\Sigma_0 \vdash \varphi$  for some finite subset  $\Sigma_0$  of  $\Sigma$ .  $\blacksquare$

**Soundness theorem.** If  $\Sigma \vdash \varphi$ , then  $\Sigma \models \varphi$ .  $\blacksquare$

**Consistency lemma.** (i)  $\Sigma$  is consistent iff  $\Sigma \not\vdash \varphi$  not for some  $\mathcal{L}$ -formula  $\varphi$ . (ii)  $\Sigma$  is consistent iff every finite subset of  $\Sigma$  is so. (iii) If  $\sigma \in \text{Sent}(\mathcal{L})$ ,  $\Sigma \cup \{\sigma\}$  is consistent iff  $\Sigma \not\vdash \neg\sigma$ .  $\blacksquare$

**Generalization lemma.** If  $\varphi(v_0, \dots, v_n) \in \text{Form}(\mathcal{L})$ , then

$$\Sigma \vdash \varphi \Rightarrow \Sigma \vdash \forall v_0 \dots \forall v_n \varphi. \quad \blacksquare$$

## 4. The completeness and model existence theorems and some of their consequences.

Let  $\mathcal{L}$  be a first-order language of type  $(\lambda, J)$ . We make the following definitions.

1. An extension  $\mathcal{L}^*$  of  $\mathcal{L}$  is called a *simple extension* of  $\mathcal{L}$  if it is obtained by adding just new constant symbols.

2. Let  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  and let  $\mathcal{L}^*$  be a simple extension of  $\mathcal{L}$ . A set  $\Sigma^* \subseteq \text{Sent}(\mathcal{L}^*)$  is called an  $\mathcal{L}$ -saturated extension of  $\Sigma$  in  $\mathcal{L}^*$  if  $\Sigma \subseteq \Sigma^*$  and, for any  $\mathcal{L}$ -formula  $\varphi$  with at most one free variable  $x$ , there is a constant symbol  $c$  of  $\mathcal{L}^*$  such that  $\Sigma^* \vdash \exists x\varphi(x) \rightarrow \varphi(c)$ .

3. A set  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is *saturated* if for any  $\mathcal{L}$ -formula  $\varphi$  with at most one free variable  $x$ , there is a constant  $c$  of  $\mathcal{L}$  for which

$$\Sigma \vdash \exists x\varphi(x) \rightarrow \varphi(c).$$

If  $\Sigma$  is saturated, then clearly:

$$\Sigma \vdash \exists x\varphi(x) \Leftrightarrow \Sigma \vdash \varphi(c) \text{ for some constant } c \text{ of } \mathcal{L}.$$

Notice also that if some set of  $\mathcal{L}$ -sentences is saturated, then  $\mathcal{L}$  contains at least one constant symbol.

**Lemma 1.** Suppose that  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is consistent. Then there is a consistent  $\mathcal{L}$ -saturated extension  $\Sigma^*$  in a simple extension  $\mathcal{L}^*$  of  $\mathcal{L}$  for which  $\|\mathcal{L}^*\| = \|\mathcal{L}\|$ .

**Proof.** Let  $F$  be the set of  $\mathcal{L}$ -formulas with at most one free variable (which we shall denote by  $x$ ). For each  $\varphi \in F$  introduce a new constant symbol  $c_\varphi$  in such a way that, if  $\varphi$  and  $\psi$  are distinct formulas, then  $c_\varphi$  and  $c_\psi$  are distinct constants. In this way we obtain a simple extension  $\mathcal{L}^*$  of  $\mathcal{L}$  clearly  $\|\mathcal{L}^*\| = \|\mathcal{L}\|$ .

Now define

$$\Sigma^* = \Sigma \cup \{\exists x\varphi(x) \rightarrow \varphi(c_\varphi) : \varphi \in F\}.$$

Clearly  $\Sigma^*$  is an  $\mathcal{L}$ -saturated extension of  $\Sigma$  in  $\mathcal{L}^*$ . It remains to show that  $\Sigma^*$  is consistent.

Suppose, on the contrary, that  $\Sigma^*$  is inconsistent. Then by the consistency lemma there is a finite subset  $\{\varphi_1, \dots, \varphi_n\}$  of  $F$  such that, writing  $c_i$  for  $c_{\varphi_i}$ ,  $\Sigma \cup \{\exists x\varphi_i \rightarrow \varphi_i(c_i) : i = 1, \dots, n\}$  is inconsistent. It follows from the consistency lemma that

$$(*) \quad \Sigma \vdash \neg \bigwedge_{i=1}^n [\exists x\varphi_i \rightarrow \varphi_i(c_i)]$$

Now choose  $n$  distinct variables  $x_1, \dots, x_n$  which do not occur in the proof from  $\Sigma$  of the sentence on the right hand side of the turnstile in (\*) – and so in particular are different from  $x$ . If in this proof we change  $c_i$  at each of its occurrences to  $x_i$  for  $i = 1, \dots, n$ , we obtain a proof of the formula  $\neg \bigwedge_{i=1}^n [\exists x\varphi_i \rightarrow \varphi_i(x_i)]$  from  $\Sigma$ , whence

$$\Sigma \vdash \neg \bigwedge_{i=1}^n [\exists x\varphi_i \rightarrow \varphi_i(x_i)].$$

By the generalization lemma,

$$\Sigma \vdash \forall v_1 \dots \forall v_n \neg \bigwedge_{i=1}^n [\exists x\varphi_i \rightarrow \varphi_i(x_i)]$$

so that

$$(**) \quad \Sigma \vdash \neg \exists v_1 \dots \exists v_n \bigwedge_{i=1}^n [\exists x\varphi_i \rightarrow \varphi_i(x_i)].$$

Now the  $x_i$  have been chosen in such a way that, if  $i \neq j$ , then  $x_i$  does not occur in  $\varphi_j(x_i)$ . So it follows from the quantifier lemma that the existential quantifiers on the right hand side of the turnstile in (\*\*) may be moved across the conjunctions and implications to yield

$$\Sigma \vdash \neg \bigwedge_{i=1}^n [\exists x \varphi_i \rightarrow \exists x_i \varphi_i(x_i)].$$

But since, clearly,  $\vdash \exists x \varphi_i \rightarrow \exists x_i \varphi_i(x_i)$  for each  $i$ , it follows that  $\Sigma$  is inconsistent, contradicting assumption. Accordingly  $\Sigma^*$  is consistent and the lemma is proved. ■

A set  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is said to be *complete* if, for any  $\sigma \in \text{Sent}(\mathcal{L})$ , we have  $\Sigma \vdash \sigma$  or  $\Sigma \vdash \neg\sigma$ .

**Lemma 2.** Suppose that  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is consistent. Then there is a complete consistent set  $\Sigma' \subseteq \text{Sent}(\mathcal{L})$  such that  $\Sigma \subseteq \Sigma'$ .

**Proof.** The family of consistent sets of sentences of  $\mathcal{L}$  containing  $\Sigma$ , ordered by inclusion, is easily seen to be closed under unions of chains, and so by Zorn's lemma has a maximal member  $\Sigma'$ . If  $\sigma \in \text{Sent}(\mathcal{L})$  and  $\Sigma' \not\vdash \sigma$ , then  $\Sigma' \cup \{\neg\sigma\}$  is consistent by the consistency lemma. Since  $\Sigma'$  is maximal consistent, we must have  $\Sigma' \cup \{\neg\sigma\} = \Sigma'$ , so *a fortiori*  $\Sigma' \vdash \neg\sigma$ . Thus  $\Sigma'$  is complete and meets the requirements of the lemma. ■

**Theorem 1.** Suppose that  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  is consistent. Then there is a simple extension  $\mathcal{L}^+$  of  $\mathcal{L}$  such that  $\|\mathcal{L}^+\| = \|\mathcal{L}\|$  and a complete saturated consistent set  $\Sigma^+ \subseteq \text{Sent}(\mathcal{L}^+)$  such that  $\Sigma \subseteq \Sigma^+$ .

**Proof.** We construct a sequence  $\mathcal{L}_0, \mathcal{L}_1, \dots$  of simple extensions of  $\mathcal{L}$  and a sequence  $\Sigma_0, \Sigma_1, \dots$  of consistent sets of sentences as follows. We begin by putting  $\mathcal{L}_0 = \mathcal{L}$  and  $\Sigma_0 = \Sigma$ . Suppose now that the consistent set  $\Sigma_n \subseteq \text{Sent}(\mathcal{L}_n)$  has been defined. By Lemma 1 there is a simple extension  $\mathcal{L}_n^*$  such that  $\|\mathcal{L}_n^*\| = \|\mathcal{L}_n\|$  and a consistent  $\mathcal{L}_n$ -saturated extension  $\Sigma_n^*$  of  $\Sigma_n$  in  $\mathcal{L}_n^*$ . And by Lemma 2, there is a complete consistent extension  $\Sigma_n^{*'}$  of  $\Sigma_n$  in  $\mathcal{L}_n^*$ : clearly  $\Sigma_n^{*'}$  is  $\mathcal{L}_n$ -saturated also. We set  $\mathcal{L}_{n+1} = \mathcal{L}_n^*$ ,  $\Sigma_{n+1} = \Sigma_n^{*'}$ . Then  $\Sigma_{n+1}$  is a complete, consistent  $\mathcal{L}_n$ -saturated extension of  $\Sigma_n$  in  $\mathcal{L}_{n+1}$ .

Now we define  $\mathcal{L}^+$  to be the union of all the languages  $\mathcal{L}_n$  and  $\Sigma^+$  to be the union of all the sets  $\Sigma_n$ . Since  $\|\mathcal{L}_n\| = \|\mathcal{L}_0\| = \|\mathcal{L}\|$  for all  $n$ , it follows that  $\|\mathcal{L}^+\| = \|\mathcal{L}\|$ . Also,  $\Sigma^+ \subseteq \text{Sent}(\mathcal{L}^+)$ ,  $\Sigma \subseteq \Sigma^+$  and  $\Sigma^+$ , as the union of the chain  $\Sigma_0 \subseteq \Sigma_1 \subseteq \dots$  of consistent sets, is itself consistent. For if  $\Sigma^+$  is inconsistent, let  $\Phi$  be the finite set of formulas of  $\mathcal{L}^+$  in a proof  $\mathcal{P}$  of a formula of the form  $\varphi \wedge \neg\varphi$  from  $\Sigma^+$ . Then  $\Phi \subseteq \text{Form}(\mathcal{L}_m)$  for some  $m$ , and  $\Sigma^+ \cap \Phi \subseteq \Sigma_n$  for some  $n$ . Writing  $q$  for the larger of  $m, n$ ,  $\mathcal{P}$  is then a proof of  $\varphi \wedge \neg\varphi$  from  $\Sigma_q$  in  $\mathcal{L}_q$ , contradicting the consistency of  $\Sigma_q$ .

Moreover,  $\Sigma^+$  is complete. for, if  $\sigma \in \text{Sent}(\mathcal{L}^+)$ , then  $\sigma \in \text{Sent}(\mathcal{L}_n)$ , for some  $n$ , and so, since  $\Sigma_n$  is complete, either  $\Sigma_n \vdash \sigma$  or  $\Sigma_n \vdash \neg\sigma$ . Since  $\Sigma_n \subseteq \Sigma^+$ , it follows that  $\Sigma^+ \vdash \sigma$  or  $\Sigma^+ \vdash \neg\sigma$ , proving the claim.

Finally,  $\Sigma^+$  is saturated. For let  $\varphi(x)$  be a formula of  $\mathcal{L}^+$  with one free variable  $x$ . Then  $\varphi(x) \in \text{Form}(\mathcal{L}_n)$  for some  $n$ . Since  $\Sigma_{n+1}$  is an  $\mathcal{L}_n$ -saturated extension of  $\Sigma_n$  in  $\mathcal{L}_{n+1}$ , there is a

constant symbol  $c$  of  $\mathcal{L}_{n+1}$  for which the sentence  $\exists x\varphi(x) \rightarrow \varphi(c)$  is provable from  $\Sigma_{n+1}$ , and hence also, since  $\Sigma_{n+1} \subseteq \Sigma^+$ , from  $\Sigma^+$ . Therefore the latter is saturated as claimed. ■

Now let  $\Sigma$  be a fixed consistent set of sentences of  $\mathcal{L}$ . Let  $C$  be the set of constant symbols of  $\mathcal{L}$ ; we shall assume that this set is nonempty. We define the relation  $\approx$  on  $C$  by

$$c \approx d \Leftrightarrow \Sigma \vdash c = d.$$

It is easy to verify, using the equality axioms in  $\mathcal{L}$ , that  $\approx$  is an equivalence relation. For each  $c \in C$  write  $\tilde{c}$  for the equivalence class of  $c$  with respect to  $\approx$ ; thus

$$\tilde{c} = \{d \in C : \Sigma \vdash c = d\}.$$

Let

$$C = \{\tilde{c} : c \in C\}$$

be the set of all such equivalence classes. Corresponding to each predicate symbol  $P_i$  of  $\mathcal{L}$  define the  $\lambda(i)$ -ary relation  $R_i$  on  $C$  by

$$R_i(c_1, \dots, c_{\lambda(i)}) \Leftrightarrow \Sigma \vdash P_i c_1 \dots c_{\lambda(i)}.$$

We can now frame the

**Definition.** The *canonical structure* determined by  $\Sigma$  is the  $\mathcal{L}$ -structure

$$\mathfrak{A}_\Sigma = (C, \{R_i : i \in I\}, \{c_j : j \in J\}).$$

Observe that  $\|\mathfrak{A}_\Sigma\| \leq |C|$ .

**Theorem 2.** Suppose that  $\Sigma$  is complete, consistent and saturated. Then  $\mathfrak{A}_\Sigma$  is a model of  $\Sigma$ .

**Proof.** We show that, for any  $\mathcal{L}$ -sentence  $\sigma$ ,

$$(*) \quad \mathfrak{A}_\Sigma \models \sigma \Leftrightarrow \Sigma \vdash \sigma.$$

That this holds for atomic sentences is an immediate consequence of the definition of  $\mathfrak{A}_\Sigma$ . We now argue by induction on the degree of complexity of the sentence  $\sigma$ .

Suppose then that  $n > 0$  and that (\*) holds for all sentences of degree  $< n$ . Let  $\sigma$  have degree  $n$ ; then  $\sigma$  is either a conjunction or a negation of sentences of degree  $< n$ , or an existentialization of a formula of degree  $< n$ . Verifying (\*) in the first two cases is routine (using the completeness of  $\Sigma$  in the negation case) and we omit the details. In the last case,  $\sigma$  is of the form  $\exists x\varphi(x)$ , where  $\varphi$  has degree  $< n$ . We then have

$$\begin{aligned} \mathfrak{A}_\Sigma \models \sigma &\Leftrightarrow \mathfrak{A}_\Sigma \models \exists x\varphi(x) \\ &\Leftrightarrow \mathfrak{A}_\Sigma \models \varphi[\tilde{c}] \text{ for some } c \in C \end{aligned}$$

(by constants lemma)

$$\Leftrightarrow \mathfrak{A}_\Sigma \models \varphi(c) \text{ for some } c \in C$$

(by (\*))

$$\Leftrightarrow \Sigma \vdash \varphi(c) \text{ for some } c \in C$$

(since  $\Sigma$  is saturated)

$$\Leftrightarrow \Sigma \vdash \exists x\varphi(x)$$

$$\Leftrightarrow \Sigma \vdash \sigma.$$

Therefore  $\sigma$  satisfies (\*) and the proof is complete. ■



These results have the following important corollaries.

**Model Existence Theorem** (Gödel-Henkin). Any consistent set  $\Sigma$  of first-order sentences has a model of cardinality at most  $\max(\aleph_0, |\Sigma|)$ .

**Proof.** Let  $\kappa = \max(\aleph_0, |\Sigma|)$ ; then  $\kappa = \|\mathcal{L}_\Sigma\|$  by the lemma on p. 3. By Theorem 1 we can extend  $\Sigma$  to a complete consistent saturated set of sentences  $\Phi$  in a simple extension  $\mathcal{L}'$  of  $\mathcal{L}_\Sigma$  such that  $\|\mathcal{L}'\| = \|\mathcal{L}_\Sigma\| = \kappa$ . By Theorem 2, the canonical structure  $\mathfrak{A}_\Phi$  is a model of  $\Phi$  and hence also of  $\Sigma$ . The expansion theorem implies that the  $\mathcal{L}_\Sigma$ -reduction  $\mathfrak{A}'$  of  $\mathfrak{A}_\Phi$  is a model of  $\Sigma$ , and that any  $\mathcal{L}$ -expansion  $\mathfrak{A}$  of  $\mathfrak{A}'$  is likewise. Moreover, if  $C$  is the set of constant symbols of  $\mathcal{L}'$ , then  $\|\mathfrak{A}\| = \|\mathfrak{A}_\Phi\| \leq |C| \leq \|\mathcal{L}_\Sigma\| = \kappa$ . The proof is complete. ■

**Completeness Theorem.** If  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  and  $\sigma \in \text{Sent}(\mathcal{L})$ , then

$$\Sigma \vdash \sigma \Rightarrow \Sigma \models \sigma.$$

**Proof.** If  $\Sigma \not\models \sigma$ , then, by the consistency theorem,  $\Sigma \cup \{\neg\sigma\}$  is consistent and so, by the model existence theorem, has a model  $\mathfrak{A}$ . Since  $\mathfrak{A}$  is a model of  $\Sigma$  but not of  $\sigma$ , it follows that  $\Sigma \not\vdash \sigma$ . ■

**Compactness Theorem.** A set of first-order sentences  $\Sigma$  has a model iff every finite subset of  $\Sigma$  has a model.

**Proof.** One way round is trivial. If, conversely, every finite subset of  $\Sigma$  has a model, then every finite subset of  $\Sigma$  is consistent and so  $\Sigma$  itself is consistent by the consistency lemma. Therefore  $\Sigma$  has a model by the model existence theorem. ■

**Invariance Theorem.** Provability and consistency are *invariant with respect to language*. That is, if  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  and  $\sigma \in \text{Sent}(\mathcal{L})$ , and  $\mathcal{L}^*$  is an extension of  $\mathcal{L}$ , then

$$(a) \Sigma \vdash \sigma \text{ in } \mathcal{L} \Leftrightarrow \Sigma \vdash \sigma \text{ in } \mathcal{L}^*$$

$$(b) \Sigma \text{ is consistent in } \mathcal{L} \Leftrightarrow \Sigma \text{ is consistent in } \mathcal{L}^*.$$

**Proof.** We prove (a); (b) is an immediate consequence. Clearly  $\Sigma \vdash \sigma$  in  $\mathcal{L} \Leftrightarrow \Sigma \vdash \sigma$  in  $\mathcal{L}^*$ . Conversely, if  $\Sigma \vdash \sigma$  in  $\mathcal{L}^*$ , then  $\Sigma \models \sigma$  by the completeness theorem, that is, every  $\mathcal{L}^*$ -structure which is a model of  $\Sigma$  is also a model of  $\sigma$ . If  $\mathfrak{A}$  is any  $\mathcal{L}$ -structure which is a model of  $\Sigma$ , it can be expanded to an  $\mathcal{L}^*$ -structure  $\mathfrak{A}^*$  which, by the expansion lemma, is also a model of  $\Sigma$ . Then  $\mathfrak{A}^*$  is a model of  $\sigma$ , and so, applying the expansion lemma again,  $\mathfrak{A}$ , as the  $\mathcal{L}$ -reduction of  $\mathfrak{A}^*$ , is a model of  $\sigma$ . Therefore, by the completeness theorem,  $\Sigma \vdash \sigma$  in  $\mathcal{L}$ . ■

**Löwenheim-Skolem Theorem.** If a set  $\Sigma$  of first-order sentences has an infinite model, it has a model of any cardinality  $\kappa \geq \max(\aleph_0, |\Sigma|)$ .

**Proof.** For simplicity write  $\mathcal{L}$  for  $\mathcal{L}_\Sigma$ . Let  $\mathcal{L}^*$  be the simple extension of  $\mathcal{L}$  obtained by adding a set  $\{d_j: j \in J\}$  of new constant symbols, where  $|J| = \kappa$ . Let

$$\Sigma^* = \Sigma \cup \{\neg(d_j = d_k) : j, k \in J \text{ \& } j \neq k\}.$$

If  $\Sigma_0$  is any finite subset of  $\Sigma^*$ , only finitely many sentences of the form  $\neg(d_j = d_k)$  occur in  $\Sigma_0$ ; let  $d_{j_1}, \dots, d_{j_n}$  be a list of all constant symbols occurring in such sentences in  $\Sigma_0$ . If now  $\mathfrak{A}$  is an infinite model of  $\Sigma$  (which we may take to be an  $\mathcal{L}$ -structure), choose  $n$  distinct elements  $a_1, \dots, a_n$  of its domain  $A$ . Let  $\mathfrak{A}^*$  be the  $\mathcal{L}$ -expansion of  $\mathfrak{A}$  in which the interpretation of  $d_{j_p}$  is  $a_p$  for  $p = 1, \dots, n$  and that of  $d_j$  is an arbitrary element of  $A$  for  $j \notin \{j_1, \dots, j_n\}$ . Clearly  $\mathfrak{A}^*$  is then a model of  $\Sigma_0$ .

It follows that every finite subset of  $\Sigma^*$  has a model. Thus every finite subset of  $\Sigma^*$  is consistent and so  $\Sigma^*$  is itself consistent. Clearly  $|\Sigma^*| = \kappa$ , so the model existence theorem implies that  $\Sigma^*$  has a model of cardinality  $\leq \kappa$ . Since the interpretations of the  $d_j$  in any model of  $\Sigma^*$  must be distinct, any such model must have cardinality  $\geq \kappa$ . So  $\Sigma^*$  has a model of cardinality  $\kappa$ ; its  $\mathcal{L}$ -reduction is a model of  $\Sigma$  of cardinality  $\kappa$ . ■

**Overspill Theorem.** If a set of first-order sentences has arbitrarily large finite models, it has an infinite model.

**Proof.** For each  $n \in \omega$  let  $\sigma_n$  be a sentence (formulable in any first-order language with equality) asserting that there are at least  $n$  individuals. Given a set  $\Sigma$  of first-order sentences, let  $\Sigma^* = \Sigma \cup \{\sigma_n : n \in \omega\}$ . If  $\Sigma$  has arbitrarily large finite models, then each finite subset of  $\Sigma^*$  has a model, so by the compactness theorem  $\Sigma^*$  has a model, which must evidently be an infinite model of  $\Sigma$ . ■

## 5. Relations between structures.

Let  $\mathfrak{A} = (A, \{R_i : i \in I\}, \{e_j : j \in J\})$  and  $\mathfrak{B} = (B, \{S_i : i \in I\}, \{d_j : j \in J\})$  be structures of the same type  $(\lambda, J)$ . We say that  $\mathfrak{A}$  is a *substructure* of  $\mathfrak{B}$ , written  $\mathfrak{A} \subseteq \mathfrak{B}$ , if  $A \subseteq B$ ,  $e_j = d_j$  for all  $j \in J$ , and  $R_i = S_i \cap A^{\lambda(i)}$  for all  $i \in I$ . If  $C$  is a nonempty subset of  $B$  containing all the designated elements of  $\mathfrak{B}$ , we define the substructure  $\mathfrak{B}|C$  of  $\mathfrak{B}$  by

$$\mathfrak{B}|C = (C, \{S_i \cap C^{\lambda(i)} : i \in I\}, \{d_j : j \in J\}).$$

An *embedding* of a structure  $\mathfrak{A}$  into a structure  $\mathfrak{B}$  is an injective map  $f : A \rightarrow B$  such that  $f(e_j) = d_j$  for all  $j \in J$ , and for all  $i \in I$  and  $a_1, \dots, a_{\lambda(i)} \in A$ , we have

$$R_i(a_1, \dots, a_{\lambda(i)}) \Leftrightarrow S_i(fa_1, \dots, fa_{\lambda(i)}).$$

If there exists an embedding of  $\mathfrak{A}$  into  $\mathfrak{B}$ , we say that  $\mathfrak{A}$  is *embeddable* into  $\mathfrak{B}$  and write  $\mathfrak{A} \subseteq \mathfrak{B}$ .

If  $f$  is an embedding of  $\mathfrak{A}$  into  $\mathfrak{B}$ , we write  $f[\mathfrak{A}]$  for the structure  $\mathfrak{B}|f[A]$ . A surjective embedding is called an *isomorphism*. If there exists an isomorphism between  $\mathfrak{A}$  and  $\mathfrak{B}$ , they are said to be *isomorphic* and we write  $\mathfrak{A} \cong \mathfrak{B}$ .

Let  $\mathcal{L}$  be the first-order language of type  $(\lambda, J)$ . We say that the  $\mathcal{L}$ -structures  $\mathfrak{A}$  and  $\mathfrak{B}$  are *elementarily equivalent*, and write  $\mathfrak{A} \equiv \mathfrak{B}$ , if  $\mathfrak{A} \models \sigma \Leftrightarrow \mathfrak{B} \models \sigma$  for any  $\mathcal{L}$ -sentence  $\sigma$ . It is easily shown that isomorphic structures are elementarily equivalent, but the Löwenheim-Skolem theorem implies that the converse fails.

The  $\mathcal{L}$ -structure  $\mathfrak{A}$  is said to be an *elementary substructure* of the  $\mathcal{L}$ -structure  $\mathfrak{B}$ , and  $\mathfrak{B}$  an *elementary extension* of  $\mathfrak{A}$ , if  $\mathfrak{A} \subseteq \mathfrak{B}$  and, for any  $\mathcal{L}$ -formula  $\varphi(v_0, \dots, v_n)$  and any  $a_0, \dots, a_n \in A$ ,

we have

$$\mathfrak{A} \models \varphi[a_0, \dots, a_n] \Leftrightarrow \mathfrak{B} \models \varphi[a_0, \dots, a_n].$$

In this situation we write  $\mathfrak{A} < \mathfrak{B}$ . Evidently  $\mathfrak{A} < \mathfrak{B} \Rightarrow \mathfrak{A} \equiv \mathfrak{B}$ , but the converse is easily seen to be false.

An embedding  $f$  of  $\mathfrak{A}$  into  $\mathfrak{B}$  is called an *elementary embedding* if for any  $\mathcal{L}$ -formula  $\varphi(v_0, \dots, v_n)$  and any  $a_0, \dots, a_n \in A$  we have

$$\mathfrak{A} \models \varphi[a_0, \dots, a_n] \Leftrightarrow \mathfrak{B} \models [fa_0, \dots, fa_n].$$

In this situation we write  $f: \mathfrak{A} < \mathfrak{B}$ . If such an  $f$  exists, we write  $\mathfrak{A} \lesssim \mathfrak{B}$ . Clearly  $\mathfrak{A} \lesssim \mathfrak{B} \Rightarrow \mathfrak{A} \equiv \mathfrak{B}$ . It is also easily shown that *any isomorphism is an elementary embedding*.

**Tarski-Vaught Lemma.** If  $\mathfrak{A}$  and  $\mathfrak{B}$  are  $\mathcal{L}$ -structures, then  $\mathfrak{A} < \mathfrak{B}$  iff  $\mathfrak{A} \subseteq \mathfrak{B}$  and, for any  $\mathcal{L}$ -formula  $\varphi(v_0, \dots, v_n)$  and any  $a_0, \dots, a_{n-1} \in A$ ,

$$(*) \quad \text{if } \mathfrak{B} \models \exists v_n \varphi[a_0, \dots, a_{n-1}], \text{ then, for some } a \in A, \mathfrak{A} \models \varphi[a_0, \dots, a_{n-1}, a].$$

**Proof.** One direction is trivial. Conversely, suppose that  $(*)$  holds. We prove by induction on the degree of  $\varphi$  that, for any  $n$ , any  $\mathcal{L}$ -formula  $\varphi(v_0, \dots, v_n)$  and any  $a_0, \dots, a_n \in A$ ,

$$(**) \quad \mathfrak{A} \models \varphi[a_0, \dots, a_n] \Leftrightarrow \mathfrak{B} \models \varphi[a_0, \dots, a_n].$$

That  $(**)$  holds for atomic formulas is obvious, as are the induction steps for  $\neg$  and  $\wedge$ . It remains to show that, if it holds for  $\varphi$ , it also holds for  $\exists v_k \varphi$ . Without loss of generality we may assume that  $n$  is greater than the index of every variable (free or bound) occurring in  $\varphi$ , and then, by making a suitable change of variable in  $\varphi$  (i.e., by substituting  $v_n$  for  $v_k$ ), that  $k = n$ .

If  $\mathfrak{A} \models \exists v_n \varphi[a_0, \dots, a_{n-1}]$ , then  $\mathfrak{A} \models \varphi[a_0, \dots, a_{n-1}, a]$  for some  $a \in A$ , and it follows from  $(**)$  for  $\varphi$  that  $\mathfrak{B} \models \varphi[a_0, \dots, a_{n-1}, a]$ , whence  $\mathfrak{B} \models \exists v_n \varphi[a_0, \dots, a_{n-1}]$ . Conversely, if  $\mathfrak{B} \models \exists v_n \varphi[a_0, \dots, a_{n-1}]$ , then, by  $(*)$ ,  $\mathfrak{A} \models \varphi[a_0, \dots, a_{n-1}, a]$  for some  $a \in A$ , so that  $\mathfrak{A} \models \exists v_n \varphi[a_0, \dots, a_{n-1}]$ . This completes the induction step and the proof. ■

**Corollary.** Write  $\mathbf{Q}$  and  $\mathbb{R}$  for the sets of rational and real numbers. Then

$$(\mathbf{Q}, \leq) < (\mathbb{R}, \leq).$$

**Proof.** We show that the Tarski-Vaught lemma applies. Suppose that, for a formula  $\varphi(v_0, \dots, v_n)$  of the appropriate language, and  $a_0 < \dots < a_{n-1} \in \mathbf{Q}$ , we have  $(\mathbb{R}, \leq) \models \exists v_n \varphi[a_0, \dots, a_{n-1}]$ . Then there is  $b \in \mathbb{R}$  such that  $(\mathbb{R}, \leq) \models \varphi[a_0, \dots, a_{n-1}, b]$ . Say  $a_i < b < a_{i+1}$  (the cases  $b <$  or  $>$  all  $a_i$  being similar). Choose  $a$  to be any rational such that  $a_i < a < a_{i+1}$ . It is easy to construct an isomorphism  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(a_j) = a_j$  for  $0 \leq j \leq n-1$  and  $f(b) = a$ . This  $f$  is also an elementary embedding. Hence  $(\mathbb{R}, \leq) \models \varphi[fa_0, \dots, fa_{n-1}, b]$ , i.e.  $(\mathbb{R}, \leq) \models \varphi[a_0, \dots, a_{n-1}, a]$ . Since  $a \in \mathbf{Q}$ , the Tarski-Vaught lemma applies to yield the required conclusion. ■

Given a set  $X$ , let  $\mathcal{L}_X$  be the simple extension of  $\mathcal{L}$  obtained by adding a set  $\{c_x: x \in X\}$  of

distinct new constant symbols indexed by  $X$ . If  $\mathfrak{A}$  is an  $\mathcal{L}$ -structure and  $X$  is a subset of its domain  $A$ , we write  $(\mathfrak{A}, X)$  for the  $\mathcal{L}_X$ -expansion of  $\mathfrak{A}$  in which the interpretation of each  $c_x$  is  $x$ . If  $f$  is a mapping of  $X$  into the domain  $B$  of an  $\mathcal{L}$ -structure  $\mathfrak{B}$ , we write  $(\mathfrak{B}, f[X])$  for the  $\mathcal{L}_X$ -expansion of  $\mathfrak{B}$  in which the interpretation of each  $c_x$  is  $f(x)$ .

The *diagram* of  $\mathfrak{A}$ ,  $\Delta(\mathfrak{A})$ , is the set of atomic and negated atomic sentences that hold in  $(\mathfrak{A}, A)$ . The *complete diagram* of  $\mathfrak{A}$ ,  $\Gamma(\mathfrak{A})$ , is the set of all sentences of  $\mathcal{L}_A$  that hold in  $(\mathfrak{A}, A)$ . The proof of the following lemma is then straightforward.

**Diagram lemma.** Let  $\mathfrak{A}$  and  $\mathfrak{B}$  be  $\mathcal{L}$ -structures. Then:

- (i)  $\mathfrak{A} \sqsubseteq \mathfrak{B}$  iff  $\mathfrak{B}$  can be expanded to a model of  $\Delta(\mathfrak{A})$ ;
- (ii)  $\mathfrak{A} \preceq \mathfrak{B}$  iff  $\mathfrak{B}$  can be expanded to a model of  $\Gamma(\mathfrak{A})$ ;
- (iii) if  $\mathfrak{A} \sqsubseteq \mathfrak{B}$ , then  $\mathfrak{A} < \mathfrak{B}$  iff  $(\mathfrak{B}, A) \models \Gamma(\mathfrak{A})$ ;
- (iv) an embedding  $f$  of  $\mathfrak{A}$  into  $\mathfrak{B}$  is an elementary embedding iff  $(\mathfrak{A}, A) \equiv (\mathfrak{B}, f[A])$ . ■

We now show that infinite structures have elementary substructures and extensions of most cardinalities.

**Theorem.** Let  $\mathfrak{A}$  be an infinite  $\mathcal{L}$ -structure.

- (i) If  $X \subseteq A$ , then for any cardinal satisfying  $\max(|X|, \|\mathcal{L}\|) \leq \kappa \leq |A|$ , there is an elementary substructure  $\mathfrak{B}$  of  $\mathfrak{A}$  such that  $|B| = \kappa$  and  $X \subseteq B$ .
- (ii)  $\mathfrak{A}$  has an elementary extension of any cardinality  $\geq \max(|X|, \|\mathcal{L}\|)$ .

**Proof.** (i) Let  $<$  be some fixed well-ordering of  $A$ . We define a sequence  $B_0, B_1, \dots$  of subsets of  $A$  recursively as follows. Choose  $B_0$  to be any subset of  $A$  such that  $|B_0| = \kappa$  and  $X \subseteq B_0$ . If  $B_n$  has been defined, put

$$B_{n+1} = \{b: \text{for some } \mathcal{L}\text{-formula } \varphi(v_0, \dots, v_m) \text{ and some } b_0, \dots, b_{m-1} \in B_n, b \text{ is the } <\text{-least element of } A \text{ such that } \mathfrak{A} \models \varphi[b_0, \dots, b_{m-1}, b]\}.$$

It is easy to check that  $B_n \subseteq B_{n+1}$  and that  $|B_{n+1}| = \kappa$ . Now define  $B$  to be the union of the  $B_n$  and  $\mathfrak{B} = \mathfrak{A} \upharpoonright B$ . Then  $\mathfrak{B}$  is a substructure of  $\mathfrak{A}$  of cardinality  $\kappa$  and it is easy to apply the Tarski-Vaught lemma to conclude that  $\mathfrak{B} < \mathfrak{A}$ .

(ii) Let  $\Gamma$  be the complete diagram of  $\mathfrak{A}$ . Then  $|\Gamma| = \max(|X|, \|\mathcal{L}\|)$ . Since  $\Gamma$  is evidently consistent, the model existence theorem implies that it has a model of any cardinality  $\kappa \geq |\Gamma| = \max(|X|, \|\mathcal{L}\|)$ . The result now follows from the diagram lemma. ■

## 6. Ultraproducts

A *filter* over a set  $I$  is a family  $\mathcal{F}$  of subsets of  $I$  such that (i)  $X, Y \in \mathcal{F} \Leftrightarrow X \cap Y \in \mathcal{F}$ , (ii)  $\emptyset \notin \mathcal{F}$ . It follows immediately from (i) that any filter  $\mathcal{F}$  over  $I$  satisfies;  $X \in \mathcal{F}$  and  $X \subseteq Y \in \mathcal{F} \Rightarrow Y \in \mathcal{F}$ . An *ultrafilter* over  $I$  is a filter  $\mathcal{U}$  over  $I$  satisfying the condition: for any  $X \in \mathcal{U}$ , either  $X \in \mathcal{U}$  or  $I - X \in \mathcal{U}$ . In particular, for any  $i \in I$ ,  $\mathcal{U}_i = \{X \subseteq I : i \in X\}$  is an ultrafilter over  $I$  called the *principal* ultrafilter generated by  $i$ . It is easily shown that an ultrafilter is precisely a filter that is maximal in the sense that it is included in no filter apart from itself. A straightforward application of Zorn's Lemma shows that a family  $\mathcal{A}$  of subsets of  $I$  is included in an ultrafilter over  $I$  if and only if it has the *finite intersection property*: that is, for any finite subfamily  $\mathcal{B}$  of  $\mathcal{A}$  we have  $\bigcap \mathcal{B} \neq \emptyset$ .

For ease of exposition we confine our attention throughout this section to structures consisting of a nonempty set and a single binary relation on that set. The appropriate language  $\mathcal{L}$  for such structures thus has a single predicate symbol of degree 2, say  $P_0$ . The type of these structures, and of  $\mathcal{L}$ , is then  $((0, 2), \emptyset)$ . It should be clear that everything we do can be extended to arbitrary structures merely by complicating the notation.

Now let  $I$  be some arbitrary fixed index set, and for each  $i \in I$  let  $\mathfrak{A}_i = (A_i, R_i)$  be an  $\mathcal{L}$ -structure. Let  $\Pi A_i$  be the Cartesian product of the sets  $A_i$ : we use letters  $f, g, h, f', g', h'$  to denote elements of  $\Pi A_i$ .

Given a family  $\mathcal{F}$  of subsets of  $I$ , we define the relation  $\sim_{\mathcal{F}}$  on  $\Pi A_i$  by

$$f \sim_{\mathcal{F}} g \Leftrightarrow \{i \in I : f(i) = g(i)\} \in \mathcal{F}.$$

It is easily shown that, if  $\mathcal{F}$  is a filter over  $I$ , then  $\sim_{\mathcal{F}}$  is an equivalence relation on  $\Pi A_i$ . From here on we shall suppose that  $\mathcal{F}$  is a filter over  $I$ . For each  $f \in \Pi A_i$  we write  $f / \mathcal{F}$  for the  $\sim_{\mathcal{F}}$ -equivalence class of  $f$ , and we define

$$\Pi A_i / \mathcal{F} = \{f / \mathcal{F} : f \in \Pi A_i\}.$$

We define the relation  $R$  on  $\Pi A_i$  by:

$$(f, g) \in R \Leftrightarrow \{i \in I : (f(i), g(i)) \in R_i\} \in \mathcal{F}.$$

It is not difficult to show that  $R$  is compatible with  $\sim_{\mathcal{F}}$  in the sense that, if  $f \sim_{\mathcal{F}} f'$  and  $g \sim_{\mathcal{F}} g'$ , then  $f R g \Rightarrow f' R g'$ . That being the case, the relation  $R$  on  $\Pi A_i$  induces the relation  $R_{\mathcal{F}}$  on  $\Pi A_i / \mathcal{F}$  given by

$$(f / \mathcal{F}, g / \mathcal{F}) \in R_{\mathcal{F}} \Leftrightarrow f R g.$$

The  $\mathcal{L}$ -structure  $\Pi \mathfrak{A}_i / \mathcal{F} = (\Pi A_i / \mathcal{F}, R_{\mathcal{F}})$  is called the *reduced product* of the family  $\{\mathfrak{A}_i : i \in I\}$  over the filter  $\mathcal{F}$ : if  $\mathcal{F}$  is an ultrafilter, the reduced product over  $\mathcal{F}$  is called an *ultraproduct*. If, for each  $i \in I$ ,  $\mathfrak{A}_i$  is a fixed structure  $\mathfrak{A}$ , the reduced product is denoted by  $\mathfrak{A} / \mathcal{F}$  and is called the *reduced power* of  $\mathfrak{A}$  over  $\mathcal{F}$ . When  $\mathcal{F}$  is an ultrafilter the reduced power is called an *ultrapower*.

Observe that if  $\mathcal{F}$  is the filter  $\{I\}$ , the reduced power  $\Pi \mathfrak{A}_i / \mathcal{F}$  is isomorphic to  $(\Pi A_i, R)$ , and that, for  $k \in I$ , the ultrapower  $\Pi \mathfrak{A}_i / \mathcal{U}_k$  is isomorphic to  $\mathfrak{A}_k$ .

If  $f = (f_0, f_1, \dots)$  is a sequence of elements of  $\Pi A_i$ , that is, if  $f \in (\Pi A_i)^{\omega}$ , we write  $f(i)$  for the sequence  $(f_0(i), f_1(i), \dots) \in A_i^{\omega}$  and, if  $\mathcal{U}$  is an ultrafilter over  $I$ ,  $f / \mathcal{U}$  for the sequence

$(f_0/\mathcal{U}, f_1/\mathcal{U}, \dots) \in (\prod A_i/\mathcal{U})^\omega$ .

We now prove the fundamental theorem on ultraproducts, viz.,

**Łoś's Theorem.** If  $\mathcal{U}$  is an ultrafilter over  $I$ ,  $\varphi$  a formula of  $\mathcal{L}$  and  $f$  a sequence of elements of  $\prod A_i$ , then

$$(*) \quad \prod \mathfrak{A}_i / \mathcal{U} \models_{f/\mathcal{U}} \varphi \Leftrightarrow \{i \in I: \mathfrak{A}_i \models_{f(i)} \varphi\} \in \mathcal{U}.$$

**Proof.** The proof goes by induction on the complexity of  $\varphi$ . That (\*) holds for atomic  $\varphi$  is a straightforward consequence of the definitions of  $\sim_{\mathcal{U}}$  and  $R_{\mathcal{U}}$ . The induction steps for  $\wedge$  and  $\neg$  follow easily from the defining properties of ultrafilters. Now suppose that (\*) holds for  $\varphi$  (and arbitrary  $f$ ); we show that it holds for  $\exists v_n \varphi$ .

Define

$$D = \{i \in I: \mathfrak{A}_i \models_{f(i)} \exists v_n \varphi\}.$$

We have to show that

$$\prod \mathfrak{A}_i / \mathcal{U} \models_{f/\mathcal{U}} \exists v_n \varphi \Leftrightarrow D \in \mathcal{U}.$$

Suppose that  $\prod \mathfrak{A}_i / \mathcal{U} \models_{f/\mathcal{U}} \exists v_n \varphi$ . Then there is some  $b \in \prod A_i$  for which  $\prod \mathfrak{A}_i / \mathcal{U} \models_{[n/b]f/\mathcal{U}} \varphi$ . Let  $E = \{i \in I: \mathfrak{A}_i \models_{([n/b]f)(i)} \varphi\}$ . Then by the induction hypothesis  $E \in \mathcal{U}$ . And since  $([n/b]f)(i) = [n/b(i)]f(i)$ , it follows that  $E \subseteq D$ , and so because  $\mathcal{U}$  is a filter,  $D \in \mathcal{U}$ .

Conversely suppose that  $D \in \mathcal{U}$ . If  $i \in D$ , then there is some  $b_i \in A_i$  such that  $\mathfrak{A}_i \models_{[n/b_i]f(i)} \varphi$ . By the axiom of choice there is  $c \in \prod A_i$  for which  $c(i) = b_i$  for every  $i \in D$ , and is an arbitrary element of  $A_i$  otherwise. Defining

$$C = \{i \in I: \mathfrak{A}_i \models_{([n/c]f)(i)} \varphi\},$$

we have  $D \subseteq C$  so that  $C \in \mathcal{U}$ . It now follows from the induction hypothesis that

$$\prod \mathfrak{A}_i / \mathcal{U} \models_{([n/c]f)/\mathcal{U}} \varphi,$$

i.e., since  $([n/c]f)/\mathcal{U} = [n/c/\mathcal{U}]f/\mathcal{U}$ ,

$$\prod \mathfrak{A}_i / \mathcal{U} \models_{[n/c/\mathcal{U}]f/\mathcal{U}} \varphi.$$

Therefore

$$\prod \mathfrak{A}_i / \mathcal{U} \models_{f/\mathcal{U}} \exists v_n \varphi,$$

completing the proof of the theorem. ■

As an immediate consequence we have the

**Corollary.** For any  $\mathcal{L}$ -sentence  $\sigma$  we have

$$\prod \mathfrak{A}_i / \mathcal{U} \models \sigma \Leftrightarrow \{i \in I: \mathfrak{A}_i \models \sigma\} \in \mathcal{U}. \quad \blacksquare$$

Let  $\mathfrak{A}$  be a structure and let  $\mathcal{U}$  be an ultrafilter on the set  $I$ . For each  $a \in A$  let  $a \in A^I$  be the function given by  $a(i) = a$  for all  $i \in I$ . The *canonical embedding* of  $\mathfrak{A}$  into  $\mathfrak{A}^I/\mathcal{U}$  is the map  $d: A \rightarrow A^I/\mathcal{U}$  defined by  $d(a) = a/\mathcal{U}$ . It is a straightforward consequence of Łoś's theorem that  $d$  is an elementary embedding.

Łoś's theorem may also be used to provide a simple direct proof of the compactness

theorem, avoiding the use of the completeness theorem. To wit, suppose that each finite subset  $\Delta$  of a given set  $\Sigma$  of sentences has a model  $\mathfrak{A}_\Delta$ ; for simplicity write  $I$  for the family of all finite subsets of  $\Sigma$ . For each  $\Delta \in I$  let  $\Delta = \{\Phi \in I : \Delta \subseteq \Phi\}$ . For any members  $\Delta_1, \dots, \Delta_n$  of  $I$ , we have

$$\Delta_1 \cup \dots \cup \Delta_n \in \Delta_1 \cap \dots \cap \Delta_n \quad ,$$

and so the collection  $\{\Delta : \Delta \in I\}$  has the finite intersection property. It can therefore be extended to an ultrafilter  $\mathcal{U}$  over  $I$ . The ultraproduct  $\prod_{\Delta \in I} \mathfrak{A}_\Delta / \mathcal{U}$  is then a model of  $\Sigma$ . For if

$\sigma \in \Sigma$ , then  $\{\sigma\} \in \Delta$ , and  $\mathfrak{A}_{\{\sigma\}} \models \sigma$ ; moreover,  $\mathfrak{A}_\Delta \models \sigma$  whenever  $\sigma \in \Delta$ . Hence

$$\{\sigma\} = \{\Delta \in I : \sigma \in \Delta\} \subseteq \{\Delta \in I : \mathfrak{A}_\Delta \models \sigma\}.$$

Since  $\{\sigma\} \in \mathcal{U}$ ,  $\{\Delta \in I : \mathfrak{A}_\Delta \models \sigma\} \in \mathcal{U}$  and therefore, by Łoś's theorem,  $\prod_{\Delta \in I} \mathfrak{A}_\Delta / \mathcal{U} \models \sigma$ . The proof is complete.

## 7. Completeness and categoricity

For simplicity, throughout this section we let  $\mathcal{L}$  be a *countable* first-order language. By a *theory* in  $\mathcal{L}$  we shall mean a set  $\Sigma$  of  $\mathcal{L}$ -sentences which is closed under provability, i.e. such that, for each  $\mathcal{L}$ -sentence  $\sigma$ , if  $\Sigma \vdash \sigma$ , then  $\sigma \in \Sigma$ . A subset  $\Gamma$  of a theory  $\Sigma$  is called a *set of postulates* for  $\Sigma$  if  $\Gamma \vdash \sigma$  for every  $\sigma \in \Sigma$ . Clearly each set  $\Gamma$  of  $\mathcal{L}$ -sentences is a set of postulates for a unique theory  $\Sigma$ , namely  $\Sigma = \{\sigma \in \text{Sent}(\mathcal{L}) : \Gamma \vdash \sigma\}$ . For each  $\mathcal{L}$ -structure  $\mathfrak{A}$  let  $\Theta(\mathfrak{A})$ , the *theory* of  $\mathfrak{A}$ , be the set of all  $\mathcal{L}$ -sentences holding in  $\mathfrak{A}$ . Clearly  $\Theta(\mathfrak{A})$  is a complete theory.

The following lemma is a straightforward consequence of the completeness theorem.

**Lemma.** The following conditions on a consistent theory  $\Sigma$  in  $\mathcal{L}$  are equivalent:

- (i)  $\Sigma$  is complete;
- (ii) any pair of models of  $\Sigma$  are elementarily equivalent;
- (iii)  $\Sigma = \Theta(\mathfrak{A})$  for some  $\mathcal{L}$ -structure  $\mathfrak{A}$ . ■

Let  $\kappa$  be an infinite cardinal. A theory  $\Sigma$  is said to be  $\kappa$ -*categorical* if any pair of models of  $\Sigma$  of cardinality  $\kappa$  are isomorphic.

**Examples.** (i) Let  $\mathcal{L}$  have no extralogical symbols and let  $\Sigma$  be the set of all  $\mathcal{L}$ -sentences which hold in every  $\mathcal{L}$ -structure. Then  $\Sigma$  is  $\kappa$ -*categorical for every infinite*  $\kappa$ .

(ii) Let  $\mathcal{L}$  have just one unary predicate symbol  $P$  and let  $\Sigma$  be the set of  $\mathcal{L}$ -sentences which hold in every  $\mathcal{L}$ -structure. Then  $\Sigma$  is *not*  $\kappa$ -*categorical for any infinite*  $\kappa$ .

(iii) Let  $\mathcal{L}$  be as in (ii) and for each natural number  $m$  let  $\sigma_m$  be the first-order sentence which asserts that there are at least  $m$  individuals having the property  $P$  and at least  $m$  individuals not having  $P$ . Let  $\Sigma$  be the theory with the set of all  $\sigma_m$  as postulates. Then  $\Sigma$  is  $\aleph_0$ -*categorical*

but not  $\kappa$ -categorical for any  $\kappa > \aleph_0$ .

(iv) Let  $\mathcal{L}$  be the language whose sole extralogical symbols are countably many constants  $c_0, c_1, \dots$  and let  $\Sigma$  be the theory with postulates  $\{\neg(c_m = c_n) : m \neq n\}$ . Then  $\Sigma$  is  $\kappa$ -categorical for every  $\kappa > \aleph_0$  but not  $\aleph_0$ -categorical.

One of the deepest results in model theory is *Morley's theorem* (whose proof is too difficult to be included here) which asserts that the four possibilities above are *exhaustive*, that is, if a theory in a countable language is  $\kappa$ -categorical for *some*  $\kappa > \aleph_0$ , it is  $\kappa$ -categorical for *all*  $\kappa > \aleph_0$ .

The next result provides a simple, but useful, sufficient condition for completeness.

**Theorem.** (Vaught's test.) Let  $\Sigma$  be a consistent theory with no finite models and which is  $\kappa$ -categorical for some infinite  $\kappa$ . Then  $\Sigma$  is complete.

**Proof.** If  $\Sigma$  is not complete, then there is a sentence  $\sigma$  such that neither  $\sigma$  nor  $\neg\sigma$  are provable from  $\Sigma$ . So both  $\Sigma \cup \{\sigma\}$  and  $\Sigma \cup \{\neg\sigma\}$  are consistent and hence have models, which must be infinite since  $\Sigma$  was assumed to have no finite models. Therefore, by Löwenheim-Skolem, both  $\Sigma \cup \{\sigma\}$  and  $\Sigma \cup \{\neg\sigma\}$  have models of cardinality  $\kappa$ . Since  $\sigma$  holds in one of these models but not in the other,  $\Sigma$  is not  $\kappa$ -categorical. ■

This theorem may be applied to establish the completeness of various theories.

**UDO** — the theory of *unbounded dense linear orderings* — is formulated in a language with just one binary predicate symbol  $R$  and has the following postulates (where we write  $x \neq y$  for  $\neg(x = y)$ ):

- (i)  $\forall x Rxx \wedge \forall x \forall y [Rxy \wedge Ryx \rightarrow x = y] \wedge \forall x \forall y \forall z [Rxy \wedge Ryz \rightarrow Rxz]$   
 $\wedge \forall x \forall y [Rxy \vee Ryx]$
- (ii)  $\forall x \forall y [Rxy \wedge x \neq y \rightarrow \exists z [x \neq z \wedge y \wedge z \wedge Rxz \wedge Rzy]]$
- (iii)  $\forall x \exists y \exists z [x \neq y \wedge x \neq z \wedge Ryx \wedge Rxz]$

Postulate (i) asserts that  $R$  is a linear ordering, (ii) that it is dense, and (iii) that it is unbounded below and above. Natural examples of models of **UDO** are  $(\mathbf{Q}, \leq)$  and  $(\mathbb{R}, \leq)$ .

**Theorem.** **UDO** is  $\aleph_0$ -categorical and so, by Vaught's test, complete.

**Proof.** Let  $(A, \leq)$  and  $(B, \leq)$  be denumerable models of **UDO**. Thus each is an unbounded dense linearly ordered set. Let  $A = \{a_n : n \in \omega\}$  and  $B = \{b_n : n \in \omega\}$ . We define two new sequences  $\{a_n^* : n \in \omega\}$  and  $\{b_n^* : n \in \omega\}$  as follows. First, put  $a_0^* = a_0$  and  $b_0^* = b_0$ . Now suppose  $k > 0$ ; we consider two cases.

(i)  $k = 2m$  is even. In this case we put  $a_k^* = a_m$ . If, for some  $j < k$ ,  $a_k^* = a_j^*$ , we put  $b_k^* = b_j^*$ . Otherwise we let  $b_k^*$  be some element of  $B$  bearing the same order relations to  $b_0^*, \dots, b_{k-1}^*$  as does  $a_k^*$  to  $a_0^*, \dots, a_{k-1}^*$ ; that is, for each  $j < k$ , if  $a_k^* > \text{or} < a_j^*$ , then  $b_k^* > \text{or} < b_j^*$ . Since  $(B, \leq)$  is a dense unbounded linearly ordered set, it is clear that such an element can always be found.

(ii)  $k = 2m + 1$  is odd. In this case we put  $b_k^* = b_m$ . If  $b_k^* = b_j$  for some  $j < k$ , put  $a_k^* = a_j^*$ . Otherwise we choose  $a_k^*$  to be some element of  $A$  bearing the same order relations to  $a_0^*, \dots, a_{k-1}^*$  as does  $b_k^*$  to  $b_0^*, \dots, b_{k-1}^*$ . Again such an element can always be found.



This completes our recursive definition. We now define  $h: A \rightarrow B$  by putting  $h(a_n^*) = b_n^*$  for each  $n \in \omega$ . Clearly  $h$  is an isomorphism between  $(A, \leq)$  and  $(B, \leq)$ . ■

The theory we consider next is most naturally formulated in a language with *operation symbols*: all our previous results extend naturally to theories in such languages.

The *language  $\mathcal{F}$  for fields* is a first-order language with constant symbols 0, 1 and binary operation symbols  $+, \cdot$ . The *theory  $\mathbf{FT}$  of fields* has the following postulates (where we write  $xy$  for  $x \cdot y$ ):

$$\begin{aligned} & \forall x \forall y [(x + y) + z = x + (y + z)] \\ & \forall x [x + 0 = x] \\ & \forall x \forall y [x + y = y + x] \\ & \forall x \exists y [x + y = 0] \\ & \forall x \forall y \forall z [(xy)z = x(yz)] \\ & \forall x [1x = x] \\ & \forall x \forall y [xy = yx] \\ & \forall x \forall y \forall z [(y + z) = xy + xz] \\ & \neg(0 = 1). \end{aligned}$$

For  $p \in \omega$ , write  $p1$  for  $1 + 1 + \dots + 1$  with  $p$  summands. If to the postulates of  $\mathbf{FT}$  we add the infinite set of sentences

$$\{\neg(p1 = 0): p \in \omega\},$$

we get the *theory  $\mathbf{FT}_0$  of fields of characteristic 0*. (Natural examples are the fields of rationals and reals.)

We now write  $x^n$  for the expression  $x \cdot (x \cdot (\dots \cdot (x \cdot x) \dots))$  with  $n$  factors. The infinite list of sentences, for  $n \geq 1$ ,

$$\forall x_0 \dots \forall x_n [\neg(x_n = 0) \rightarrow \exists y (x_n y^n + x_{n-1} y^{n-1} + \dots + x_1 y + x_0 = 0)]$$

when added to the postulates of  $\mathbf{FT}_0$ , yields the *theory  $\mathbf{ACF}_0$  of algebraically closed fields of characteristic 0*. Each new postulate asserts that all polynomials of a given degree  $n$  has a zero.

We observe that  $\mathbf{ACF}_0$  is *not*  $\aleph_0$ -categorical. For the field  $\mathbf{F}$  of algebraic numbers and the algebraic closure of the field  $\mathbf{F}[\pi]$  obtained by adjoining the transcendental  $\pi$  to  $\mathbf{F}$  are countable nonisomorphic models of  $\mathbf{ACF}_0$ . On the other hand, a classical theorem of Steinitz asserts that  $\mathbf{ACF}_0$  is  $\kappa$ -categorical for any *uncountable*  $\kappa$ , so we conclude from Vaught's test that  $\mathbf{ACF}_0$  is *complete*. Since the field  $\mathbb{C}$  of complex numbers is a model of  $\mathbf{ACF}_0$ , it follows that  $\mathbf{ACF}_0$  is a *set of postulates for the theory of  $\mathbb{C}$* .

## 8. The elementary chain theorem and some of its consequences.

Let  $\mathfrak{A}_0 \subseteq \mathfrak{A}_1 \subseteq \dots$  be a chain of  $\mathcal{L}$ -structures: in particular the  $\mathfrak{A}_i$  all have the same designated elements. The *union* of the chain is the structure  $\mathfrak{A} = \bigcup_{n \in \omega} \mathfrak{A}_n$  defined as follows. The

domain of  $\mathfrak{A}$  is the set  $A = \bigcup_{n \in \omega} A_n$ . For  $i \in I$ , the  $i^{\text{th}}$  relation  $R_i$  of  $\mathfrak{A}$  is the union of the corresponding  $i^{\text{th}}$  relations of the  $A_n$ . The designated elements of  $\mathfrak{A}$  are the designated elements of the  $\mathfrak{A}_n$ . Clearly each  $\mathfrak{A}_n$  is a substructure of  $\mathfrak{A}$ .

A chain of structures  $\mathfrak{A}_0 \subseteq \mathfrak{A}_1 \subseteq \dots$  in which each  $\mathfrak{A}_n$  is an elementary substructure of  $\mathfrak{A}_{n+1}$  is called an *elementary chain*. In this case we write  $\mathfrak{A}_0 < \mathfrak{A}_1 < \dots$ .

**Elementary Chain Theorem.** Each member of an elementary chain of structures is an elementary substructure of the union of the chain.

**Proof.** Let  $\mathfrak{A}_0 < \mathfrak{A}_1 < \dots$  be an elementary chain, and let  $\mathfrak{A}$  be its union. We prove the following assertion by induction on the degree of a formula: for any  $\mathcal{L}$ -formula  $\varphi(v_0, \dots, v_n)$ , any  $n \in \omega$  and any  $a_0, \dots, a_m \in A_n$ ,

$$(*) \quad \mathfrak{A}_n \models \varphi[a_0, \dots, a_m] \Leftrightarrow \mathfrak{A} \models \varphi[a_0, \dots, a_m].$$

The proof is routine for atomic formulas, and the induction steps for  $\neg$  and  $\wedge$  are easy. Now suppose that  $\varphi$  is existential; without loss of generality we may assume that  $\varphi$  is  $\exists v_n \psi$ , and that  $\psi$  satisfies (\*).

If  $a_0, \dots, a_{m-1} \in A_n$  and  $\mathfrak{A}_n \models \varphi[a_0, \dots, a_{m-1}]$ , then for some  $a \in A_n$  we have  $\mathfrak{A}_n \models \psi[a_0, \dots, a_{m-1}, a]$ . So by (\*)  $\mathfrak{A} \models \psi[a_0, \dots, a_{m-1}, a]$  whence  $\mathfrak{A} \models \varphi[a_0, \dots, a_{m-1}]$ .

Conversely, suppose that  $\mathfrak{A} \models \varphi[a_0, \dots, a_{m-1}]$ . Then  $\mathfrak{A} \models \psi[a_0, \dots, a_{m-1}, a]$  for some  $a \in A$ . For some  $k$ ,  $a \in A_k$ . Let  $\ell$  be the larger of  $k$  and  $n$ . Then  $a_0, \dots, a_{m-1}, a \in A_\ell$  and so, by (\*),  $\mathfrak{A}_\ell \models \psi[a_0, \dots, a_{m-1}, a]$ , whence  $\mathfrak{A}_\ell \models \varphi[a_0, \dots, a_{m-1}]$ . But  $n \leq \ell$  and so, since  $\mathfrak{A}_n < \mathfrak{A}_\ell$ , we conclude that  $\mathfrak{A}_n \models \varphi[a_0, \dots, a_{m-1}]$ . ■

We use this in the proof of the

**Joint Consistency Theorem.** Let  $\Sigma$  and  $\Pi$  be theories in  $\mathcal{L}$ , and let  $\mathcal{E}$  be the language whose extralogical symbols are those common to  $\mathcal{L}_\Sigma$  and  $\mathcal{L}_\Pi$ . Then the following are equivalent:

- (i)  $\Sigma \cup \Pi$  is consistent.;
- (ii) for no  $\mathcal{E}$ -sentence  $\sigma$  do we have  $\Sigma \vdash \sigma$  and  $\Pi \vdash \neg\sigma$ ;
- (iii) for some complete (consistent) theory  $\Delta$  in  $\mathcal{E}$ , both  $\Sigma \cup \Delta$  and  $\Pi \cup \Delta$  are consistent;
- (iv) there is an  $\mathcal{E}$ -structure which can be expanded both to a model of  $\Sigma$  and to a model of  $\Pi$ .

**Proof.** (i)  $\Rightarrow$  (ii) is obvious.

(ii)  $\Rightarrow$  (iii). Assume (ii) and let  $\Sigma^* = \{\sigma \in \text{Sent}(\mathcal{E}) : \Sigma \vdash \sigma\}$ . It follows easily from (ii) that  $\Pi \cup \Sigma^*$  is consistent and so has a model  $\mathfrak{A}$ . Let  $\Delta$  be the theory of the  $\mathcal{E}$ -structure  $\mathfrak{A} \upharpoonright \mathcal{E}$ . Since  $\mathfrak{A} \models \Pi \cup \Delta$ ,  $\Pi \cup \Delta$  is consistent. If  $\Sigma \cup \Delta$  is inconsistent, there is  $\sigma \in \Delta$  such that  $\Sigma \vdash \neg\sigma$ , i.e.  $\neg\sigma \in \Sigma^*$ . But then  $\mathfrak{A} \models \neg\sigma$ , whence  $\neg\sigma \in \Delta$ , a contradiction. Hence  $\Sigma \cup \Delta$  is consistent.

(iii)  $\Rightarrow$  (iv). Assume (iii), and let  $\mathfrak{A}_0$  and  $\mathfrak{B}_0$  be models of  $\Sigma \cup \Delta$  and  $\Pi \cup \Delta$ , respectively. Then since  $\mathfrak{A}_0 \upharpoonright \mathcal{E}$  and  $\mathfrak{B}_0 \upharpoonright \mathcal{E}$  are both models of the complete theory  $\Delta$ , they are elementarily equivalent. It follows easily from this that the union  $\Gamma$  of the complete diagram  $\Gamma^*$  of  $\mathfrak{A}_0 \upharpoonright \mathcal{E}$  with

the complete diagram  $\Gamma^{**}$  of  $\mathfrak{B}_0$  is consistent. (Observe that each finite subset of  $\Gamma^*$  is interpretable in  $\mathfrak{B}_0$ .) Let  $\mathfrak{B}^*$  be a model of  $\Gamma$  and let  $\mathfrak{B}_1$  be its  $\mathcal{L}$ -reduction. Then since  $\mathfrak{B}^*$  is a model of both  $\Gamma^*$  and  $\Gamma^{**}$  it follows from the diagram lemma that  $\mathfrak{A}_0 \mid \mathcal{E} \simeq \mathfrak{B}_1 \mid \mathcal{E}$  and  $\mathfrak{B}_0 \simeq \mathfrak{B}_1$ . Identifying  $\mathfrak{B}_0$  with its image in  $\mathfrak{B}_1$  makes the former an elementary substructure of the latter. Let  $f_1$  be an elementary embedding of  $\mathfrak{A}_0 \mid \mathcal{E}$  into  $\mathfrak{B}_1 \mid \mathcal{E}$ .

Passing to the extended language  $\mathcal{E}_{A_0}$ , the diagram lemma implies that the structures  $(\mathfrak{A}_0 \mid \mathcal{E}, A_0) = (\mathfrak{A}_0, A_0) \mid \mathcal{E}_{A_0}$  and  $(\mathfrak{B}_1 \mid \mathcal{E}, f_1[A_0]) = (\mathfrak{B}_1, f_1[A_0])$  are elementarily equivalent. Repeating the above construction in the other direction, this time with the  $\mathcal{L}_{A_0}$ -structures  $(\mathfrak{A}_0, A_0)$  and  $(\mathfrak{B}_1, f_1[A_0])$  in place of  $\mathfrak{A}_0, \mathfrak{B}_0$ , respectively, we obtain an elementary extension  $\mathfrak{A}_1$  of  $\mathfrak{A}_0$  and an elementary embedding  $g_1$  of  $(\mathfrak{B}_1, f_1[A_0]) \mid \mathcal{E}_{A_0}$  into  $(\mathfrak{A}_1, A_0) \mid \mathcal{E}_{A_0}$ . Then  $g \circ f_1$  is the identity on  $A_0$ , so that  $f_1 \subseteq g_1^{-1}$ .

Iterating this construction yields a diagram

$$\begin{array}{ccccccc} \mathfrak{A}_0 & < & \mathfrak{A}_1 & < & \mathfrak{A}_2 & < & \dots \\ & \searrow f_1 & \uparrow g_1 & & \searrow f_2 & \uparrow g_2 & \\ \mathfrak{B}_0 & < & \mathfrak{B}_1 & < & \mathfrak{B}_2 & < & \dots \end{array}$$

such that, for each  $m$ ,  $f_m$  is an elementary embedding of  $\mathfrak{A}_{m-1} \mid \mathcal{E}$  into  $\mathfrak{B}_m \mid \mathcal{E}$ ,  $g_m$  is an elementary embedding of  $\mathfrak{B}_m \mid \mathcal{E}$  into  $\mathfrak{A}_m \mid \mathcal{E}$ , and  $f_m \subseteq g_m^{-1} \subseteq f_{m+1}$ . Let  $\mathfrak{A}$  and  $\mathfrak{B}$  be the unions of the elementary chains  $\mathfrak{A}_0 < \mathfrak{A}_1 < \dots$  and  $\mathfrak{B}_0 < \mathfrak{B}_1 < \dots$  respectively. Then, by the elementary chain theorem,  $\mathfrak{A}$  is a model of  $\Sigma$  and  $\mathfrak{B}$  is a model of  $\Pi$ . Moreover,  $\bigcup_{m \in \mathfrak{w}} f_m$  is an isomorphism of  $\mathfrak{A} \mid \mathcal{E}$  and  $\mathfrak{B} \mid \mathcal{E}$  (since, by construction, it has inverse  $\bigcup_{m \in \mathfrak{w}} g_m$ ). It follows that  $\mathfrak{B}$  is isomorphic to a structure  $\mathfrak{B}'$  such that  $\mathfrak{A} \mid \mathcal{E} = \mathfrak{B}' \mid \mathcal{E}$ . Accordingly the  $\mathcal{E}$ -structure  $\mathfrak{A} \mid \mathcal{E}$  can be expanded both to the model  $\mathfrak{A}$  of  $\Sigma$  and to the model  $\mathfrak{B}'$  of  $\Pi$ .

(iv)  $\Rightarrow$  (i). Let  $\mathfrak{A}$  be an  $\mathcal{E}$ -structure expandable both to a model  $\mathfrak{B}$  of  $\Sigma$  and to a model  $\mathfrak{C}$  of  $\Pi$ . Define the  $\mathcal{L}$ -structure  $\mathfrak{D}$  as follows: the domain of  $\mathfrak{D}$  is that of  $\mathfrak{A}$ ; if  $s$  is any extralogical symbol of  $\mathcal{L}$ , then

$$s^{\mathfrak{D}} = \begin{cases} s^{\mathfrak{A}} & \text{if } s \in \mathcal{E} \\ s^{\mathfrak{B}} & \text{if } s \in \mathcal{L} - \mathcal{A}_\Sigma \\ s^{\mathfrak{C}} & \text{if } s \in \mathcal{A}_\Pi \end{cases}$$

Clearly  $\mathfrak{D} \mid \mathcal{A}_\Sigma = \mathfrak{B}$ , so  $\mathfrak{D} \models \Sigma$ . Also,  $\mathfrak{D} \mid \mathcal{A}_\Pi = \mathfrak{C}$ , so  $\mathfrak{D} \models \Pi$ . Therefore  $\mathfrak{D}$  is a model of  $\Sigma \cup \Pi$ , so the latter is consistent. ■

From this we deduce

**Craig's Interpolation Theorem.** Suppose  $\sigma, \tau$  are  $\mathcal{L}$ -sentences and  $\vdash \sigma \rightarrow \tau$ . Then there is a sentence  $\theta$  such that  $\vdash \sigma \rightarrow \theta$ ,  $\vdash \theta \rightarrow \tau$ , and every extralogical symbol occurring in  $\theta$  occurs in both  $\sigma$  and  $\tau$ .

**Proof.** Let  $\mathcal{L}$  be the language whose extralogical symbols are exactly those occurring in both  $\sigma$  and  $\tau$ . If  $\vdash \sigma \rightarrow \tau$ , then  $\{\sigma, \neg\tau\}$  is inconsistent, so by (ii) of the joint consistency theorem there is an  $\mathcal{L}$ -sentence  $\theta$  such that  $\sigma \vdash \theta$  and  $\neg\tau \vdash \neg\theta$ . The result now follows immediately. ■

Suppose that  $\Sigma \subseteq \text{Sent}(\mathcal{L})$  contains the  $n$ -ary predicate symbol  $P$ .  $P$  is said to be *explicitly definable* from  $\Sigma$  if there is an  $\mathcal{L}$ -formula  $\varphi(x_1, \dots, x_n)$ , in which  $P$  does not occur, such that

$$\Sigma \vdash \forall x_1 \dots \forall x_n [Px_1 \dots x_n \leftrightarrow \varphi].$$

Now let  $P^*$  be an  $n$ -ary predicate symbol *not* belonging to  $\mathcal{L}$ , and let  $\Sigma^*$  be the set of sentences obtained from  $\Sigma$  by replacing all occurrences of  $P$  by  $P^*$ . Then  $P$  is said to be *implicitly definable* from  $\Sigma$  if

$$\Sigma \cup \Sigma^* \vdash \forall x_1 \dots \forall x_n [Px_1 \dots x_n \leftrightarrow P^*x_1 \dots x_n].$$

Semantically speaking, this means that any pair of  $\mathcal{L}$ -structures which are both models of  $\Sigma$ , have the same domain and agree on the interpretation of all extralogical symbols apart possibly from  $P$ , must also agree on the interpretation of  $P$ .

Clearly, if  $P$  is explicitly definable from  $\Sigma$ , it is implicitly definable from  $\Sigma$ . Conversely, we have

**Beth's Definability Theorem.** If  $P$  is implicitly definable from  $\Sigma$ , it is explicitly definable from  $\Sigma$ .

**Proof.** Suppose  $P$  is implicitly definable from  $\Sigma$ . Without loss of generality we may assume  $\Sigma$  to be finite, and we can then replace  $\Sigma$  by the conjunction of all its sentences. So we may assume that  $\Sigma$  consists of a single sentence  $\sigma$ . Let  $\sigma^*$  be the result of replacing each occurrence of  $P$  in  $\sigma$  by  $P^*$ . Then we have

$$(1) \quad \{\sigma, \sigma^*\} \vdash \forall x_1 \dots \forall x_n [Px_1 \dots x_n \rightarrow P^*x_1 \dots x_n].$$

Now add new constant symbols  $c_1, \dots, c_n$  to  $\mathcal{L}$ . Then, by (1),

$$\{\sigma, \sigma^*\} \vdash Pc_1 \dots c_n \rightarrow P^*c_1 \dots c_n.$$

So

$$\vdash \sigma \wedge Pc_1 \dots c_n \rightarrow (\sigma^* \rightarrow P^*c_1 \dots c_n).$$

By Craig's theorem, there is a sentence  $\theta$  whose extralogical symbols are common to both  $\sigma \wedge Pc_1 \dots c_n$  and  $\sigma^* \rightarrow P^*c_1 \dots c_n$ , hence, in particular, not containing  $P$  or  $P^*$  such that  $\vdash \sigma \wedge Pc_1 \dots c_n \rightarrow \theta$  and  $\vdash \theta \rightarrow (\sigma^* \rightarrow P^*c_1 \dots c_n)$ .

Therefore

$$(2) \quad \sigma \vdash Pc_1 \dots c_n \rightarrow \theta$$

and

$$(3) \quad \sigma^* \vdash \theta \rightarrow P^*c_1 \dots c_n.$$

If we replace  $P^*$  by  $P$  in (3),  $\sigma^*$  becomes  $\sigma$  and  $\theta$  is unchanged. So

$$(4) \quad \sigma \vdash \theta \rightarrow Pc_1 \dots c_n.$$

(2) and (4) now give

$$(5) \quad \Sigma \vdash \theta \leftrightarrow Pc_1 \dots c_n.$$

But  $\theta$  is  $\varphi(c_1, \dots, c_n)$  for some  $\mathcal{L}$ -formula  $\varphi(x_1, \dots, x_n)$  in which  $P$  does not occur. Since  $c_1, \dots, c_n$

are not in  $\mathcal{L}$ , the result of replacing  $c_i$  by  $x_i$  ( $i = 1, \dots, n$ ) in the proof from  $\Sigma$  of  $\theta \leftrightarrow P_{c_1 \dots c_n}$  yields a proof from  $\Sigma$  of  $\varphi \leftrightarrow P_{x_1 \dots x_n}$ . Applying the generalization lemma gives

$$\Sigma \vdash \forall x_1 \dots \forall x_n [\varphi \leftrightarrow P_{x_1 \dots x_n}]$$

and so  $P$  is explicitly definable from  $\Sigma$ . ■